



Contents lists available at ScienceDirect

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv

Early stage litter decomposition across biomes

Ika Djukic^{a,*}, Sebastian Kepfer-Rojas^b, Inger Kappel Schmidt^b, Klaus Steenberg Larsen^b,
Claus Beier^b, Björn Berg^c, Kris Verheyen^d, TeaComposition:

Adriano Caliman¹, Alain Paquette², Alba Gutiérrez-Girón³, Alejandro Valdecantos⁴, Alessandro Petraglia⁵, Heather Alexander⁶, Algirdas Augustaitis⁷, Amélie Saillard⁸, Ana Carolina Ruiz Fernández⁹, Ana I. Sousa¹⁰, Ana I. Lillebø¹⁰, Anderson da Rocha Gripp¹¹, André-Jean Francez¹², Andrea Fischer¹³, Andreas Bohner¹⁴, Andrey Malyshev¹⁵, Andrijana Andrić¹⁶, Andy Smith¹⁷, Angela Stanisci¹⁸, Anikó Seres¹⁹, Anja Schmidt²⁰, Anna Avila²¹, Anne Probst²², Annie Ouin²², Anzar A. Khuroo²³, Arne Verstraeten²⁴, Artur Stefanski²⁵, Aurora Gaxiola²⁶, Bart Muys²⁷, Bernard Bosman²⁸, Bernd Ahrends²⁹, Bill Parker³⁰, Birgit Sattler³¹, Björn Berg³², Bo Yang^{33,34}, Bohdan Juráni³⁵, Brigitta Erschbamer³⁶, Carmen Eugenia Rodriguez Ortiz³⁷, Casper T. Christiansen³⁸, E. Carol Adair³⁹, Céline Meredieu⁴⁰, Cendrine Mony⁴², Charles Andrew Nock⁴¹, Chi-Ling Chen⁴², Chiao-Ping Wang⁴³, Christel Baum⁴⁴, Christian Rixen⁴⁵, Christine Delire⁴⁶, Christophe Piscart⁴², Christopher Andrews⁴⁷, Claus Beier⁴⁸, Corinna Rebmann⁴⁹, Cristina Branquinho⁵⁰, Dana Polyanskaya⁵¹, David Fuentes Delgado⁴, Dirk Wundram⁵², Diyaa Radeideh^{53,54}, Eduardo Ordóñez-Regil⁵⁵, Edward Crawford⁵⁶, Elena Preda⁵⁷, Elena Tropina⁵¹, Elli Groner⁵⁸, Eric Lucot⁵⁹, Erzsébet Hornung⁶⁰, Esperança Gacia⁶¹, Esther Lévesque⁶², Evanilde Benedito⁶³, Evgeny A. Davydov^{64,65}, Evy Ampoorter⁶⁶, Fabio Padilha Bolzan⁶⁷, Felipe Varela⁶⁸, Ferdinand Kristöfel⁶⁹, Fernando Maestre⁷⁰, Florence Maunoury-Danger⁷¹, Florian Hofhansl⁷², Florian Kitz⁷³, Flurin Sutter⁷⁴, Francisco Cuesta^{75,76}, Francisco de Almeida Lobo⁷⁷, Franco Leandro de Souza⁶⁷, Frank Berninger³², Franz Zehetner^{78,79}, Georg Wohlfahrt⁷³, George Vourlitis⁸⁰, Geovana Carreño-Rocabado^{81,82}, Gina Arena⁸³, Gisele Daiane Pinha⁶³, Grizelle González⁸⁴, Guylaine Canut⁴⁶, Hanna Lee³⁸, Hans Verbeeck⁸⁵, Harald Auge^{20,86}, Harald Pauli^{87,88}, Hassan Bismarck Nacro⁸⁹, Héctor Bahamonde⁹⁰, Heike Feldhaar⁹¹, Heinke Jäger⁹², Helena C. Serrano⁵⁰, Hélène Verheyden⁹³, Helge Bruelheide^{34,86}, Henning Meessenburg⁹⁴, Hermann Jungkunst⁹⁵, Hervé Jactel⁴⁰, Hideaki Shibata⁹⁶, Hiroko Kurokawa⁹⁷, Hugo López Rosas⁹⁸, Hugo L. Rojas Villalobos⁹⁹, Ian Yesilonis¹⁰⁰, Ika Djukic¹⁰¹, Inara Melece¹⁰², Inge Van Halder⁴⁰, Inger Kappel Schmidt⁴⁸, Inmaculada Garcia Quiros⁴⁹, Isaac Makelele¹⁰³, Issaka Senou¹⁰⁴, István Fekete¹⁰⁵, Ivan Mihal¹⁰⁶, Ivika Ostonen¹⁰⁷, Jana Borovská¹⁰⁸, Javier Roales¹⁰⁹, Jawad Shoqir^{53,54}, Jean-Christophe Lata¹¹⁰, Jean-Paul Theurillat^{111,112}, Jean-Luc Probst²², Jess Zimmerman¹¹³, Jeyanny Vijayanathan¹¹⁴, Jianwu Tang¹¹⁵, Jill Thompson¹¹⁶, Jiri Dolezal¹¹⁷, Joan-Albert Sanchez-Cabeza¹¹⁸, Joël Merlet⁹³, Joh Henschel¹¹⁹, Johan Neirynck²⁴, Johannes Knops¹²⁰, John Loehr¹²¹, Jonathan von Oppen⁴⁵, Jónína Sigríður Þorlákssdóttir¹²², Jörg Löffler⁵², José Gilberto Cardoso Mohedano¹²³, José-Luis Benito-Alonso¹²⁴, Jose Marcelo Torezan¹²⁵, Joseph C. Morina¹²⁶, Juan J. Jiménez¹²⁷, Juan Dario Quinde¹²⁸, Juha Alatalo¹²⁹, Julia Seeber¹³⁰, Jutta Stadler²⁰, Kaie Kriiska¹⁰⁷, Kalifa Coulibaly⁸⁹, Karibu Fukuzawa¹³², Katalin Szlavecz¹³³, Katarína Gerhátoová¹⁰⁸, Kate Lajtha¹³⁴, Kathrin Käppeler¹³⁵, Katie A. Jennings¹³⁶, Katja Tielbörger¹³⁷, Kazuhiko Hoshizaki¹³⁸, Ken Green¹³⁹, Klaus Steenberg Larsen⁴⁸, Kris Verheyen¹⁴⁰, Lambiénou Yé¹⁰⁴, Laryssa Helena Ribeiro Pazianoto⁶³, Laura Dienstbach⁴⁹, Laura Williams¹⁴¹, Laura Yahdjian¹⁴², Laurel M. Brigham¹⁴³, Liesbeth van den Brink¹⁴⁴,

* Corresponding author at: Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse 111, 8903 Birmensdorf, Zürich, Switzerland.

E-mail addresses: ika.djukic@umweltbundesamt.at (I. Djukic), skro@ign.ku.dk (S. Kepfer-Rojas), iks@ign.ku.dk (I.K. Schmidt), kstl@ign.ku.dk (K.S. Larsen), cbe@ign.ku.dk (C. Beier), Kris.Verheyen@UGent.be (K. Verheyen), algutier@ucm.es (A. Gutiérrez-Girón), a.valdecantos@ua.es (A. Valdecantos), alessandro.petraglia@unipr.it (A. Petraglia),

Lindsey Rustad¹⁴⁵, Lipeng Zhang³³, Lourdes Morillas¹⁴⁶, Luciana Silva Carneiro¹, Luciano Di Martino¹⁴⁷, Luis Villar¹²⁷, Maaikje Y. Bader¹⁴⁸, Madison Morley¹⁴³, Marc Lebourvier¹⁴⁹, Marcello Tomaselli⁵, Marcelo Sternberg¹⁵⁰, Marcus Schaub⁷⁴, Margarida Santos-Reis⁵⁰, Maria Glushkova¹⁵¹, María Guadalupe Almazán Torres⁵⁵, Marie-Andrée Giroux¹⁵², Marie-Anne de Graaff¹⁵³, Marie-Noëlle Pons¹⁵⁴, Marijn Bauters¹⁵⁵, Marina Mazón¹⁵⁶, Mark Frenzel²⁰, Markus Didion¹⁵⁷, Markus Wagner²⁹, Maroof Hamid²³, Marta L. Lopes¹⁰, Martha Apple¹⁵⁸, Martin Schädler^{20,86}, Martin Weih¹⁵⁹, Matteo Gualmini⁵, Matthew A. Vadeboncoeur¹⁶⁰, Michael Bierbaumer¹⁶¹, Michael Danger¹⁶², Michael Liddell¹⁶³, Michael Mirtl¹⁶⁴, Michael Scherer-Lorenzen⁴¹, Michal Ruzek^{165,166}, Michele Carbognani⁵, Michele Di Musciano¹⁶⁷, Michinari Matsushita¹⁶⁸, Miglena Zhiyanski¹⁵¹, Mihai Pușcaș¹⁶⁹, Milan Barna¹⁰⁶, Mioko Ataka¹⁷⁰, Mohammed Alsafran¹²⁹, Monique Carnol²⁸, Nadia Barsoum¹⁷¹, Naoko Tokuchi¹⁷², Nico Eisenhauer⁸⁶, Nicolas Lecomte¹⁷³, Nina Filippova¹⁷⁴, Norbert Hölzel¹⁷⁵, Olga Ferlian⁸⁶, Oscar Romero¹²⁸, Pablo Peri⁹⁰, Paige Weber¹⁷⁶, Pascal Vittoz¹⁷⁷, Pavel-Dan Turtureanu¹⁷⁸, Peter Fleischer¹⁷⁹, Peter Macreadie¹⁸⁰, Peter Haase^{181,182}, Peter Reich^{183,184}, Petr Petřík¹⁸⁵, Philippe Choler⁸, Pierre Marmonier¹⁸⁶, Priscilla Muriel⁶⁸, Quentin Ponette¹⁸⁷, Rafael Dettogni Guariento⁶⁷, Rafaella Canessa¹⁴⁸, Ralf Kiese¹⁸⁸, Rebecca Hewitt¹⁸⁹, Regin Rønn¹⁹⁰, Rita Adrian¹⁹¹, Róbert Kanka¹⁹², Robert Weigel¹⁹³, Roberto Cazzolla Gatti¹⁹⁴, Rodrigo Lemes Martins¹⁹⁵, Romain Georges¹², Rosa Isela Meneses¹⁹⁶, Rosario G. Gavilán³, Sabyasachi Dasgupta¹⁹⁷, Sally Wittlinger¹⁹⁸, Sara Puijalon¹⁸⁶, Sarah Freda¹⁷⁶, Satoshi Suzuki¹⁹⁹, Sean Charles²⁰⁰, Sébastien Gogo²⁰¹, Sebastian Kepfer-Rojas⁴⁸, Simon Drollinger²⁰², Simone Mereu²⁰³, Sonja Wipf⁴⁵, Stacey Trevathan-Tackett²⁰⁴, Stefan Löfgren²⁰⁵, Stefan Stoll^{95,206}, Stefan Trogisch^{34,86}, Stefanie Hoerber¹⁵⁹, Steffen Seitz¹³⁵, Stephan Glatzel²⁰², Sue J. Milton²⁰⁷, Sylvie Dousset²⁰⁸, Taiki Mori²⁰⁹, Takanori Sato²¹⁰, Takeshi Ise¹⁷², Takuo Hishi²¹¹, Tanaka Kenta²¹², Tatsuro Nakaji²¹³, Thaisa Sala Michelan²¹⁴, Thierry Camboulive²¹⁵, Thomas J. Mozdzer¹⁷⁶, Thomas Scholten¹³⁵, Thomas Spiegelberger²¹⁶, Thomas Zechmeister²¹⁷, Till Kleinebecker¹⁷⁵, Tsutomu Hiura²¹³, Tsutomu Enoki²¹⁸, Tudor-Mihai Ursu²¹⁹, Umberto Morra di Cella²²⁰, Ute Hamer¹⁷⁵, Valentin Klaus^{175,233}, Vanessa Mendes Rêgo²²¹, Valter Di Cecco¹⁴⁷, Verena Busch¹⁷⁵, Veronika Fontana¹³⁰, Veronika Piscová¹⁰⁸, Victoria Carbonell^{188,222}, Victoria Ochoa⁷⁰, Vincent Bretagnolle²²³, Vincent Maire⁶², Vinicius Farjalla²²⁴, Wenjun Zhou²²⁵, Wentao Luo²²⁶, William H. McDowell²²⁷, Yalin Hu²²⁸, Yasuhiro Utsumi²²⁹, Yuji Kominami¹⁷⁰, Yulia Zaika²³⁰, Yury Rozhkov²³¹, Zsolt Kotroczó²³², Zsolt Tóth⁶⁰

¹ Universidade Federal do Rio Grande do Norte, Departamento de Ecologia, 59078-900 Natal, RN, Brazil

² Université du Québec à Montréal, Centre for Forest Research, P.O. Box 8888, Centre-ville Station, Montreal, QC H3C 3P8, Canada

³ Dpto. Biología Vegetal II, Facultad de Farmacia, Universidad Complutense, E-28040 Madrid, Spain

⁴ CEAM Foundation (Mediterranean Center for Environmental Studies), Department of Ecology, University of Alicante, Carretera San Vicente del Raspeig s/n 03690, San Vicente del Raspeig, Alicante, Spain

⁵ Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità ambientale, Parco Area delle Scienze 11/A, I-43124 Parma, Italy

⁶ Mississippi State University, Department of Forestry, 327 Thompson Hall, 775 Stone Blvd., P.O. Box 9681, MS 39762, USA

⁷ Aleksandras Stulginskis University, Forest Monitoring Laboratory, Kaunas distr., Studentu 13, LT-53362, Lithuania

⁸ Univ. Grenoble Alpes, CNRS, LECA, LTSEZ Zone Atelier Alpes, F-38000 Grenoble, France

⁹ Unidad Académica Mazatlán, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Calz. Joel Montes Camarena s/n, 82040 Mazatlán, Sinaloa, Mexico

¹⁰ Department of Biology & CESAM, University of Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

¹¹ Universidade Federal do Rio de Janeiro (UFRJ), Departamento de Ecologia, Instituto de Biologia, CCS, Bloco A, Ilha do Fundão, Rio de Janeiro, RJ CEP: 21.941-590, Brazil

¹² ECOBIO, CNRS-Université de Rennes 1 & LTSEZ Zone Atelier Armorique, Avenue du Général Leclerc, 35042 Rennes Cedex, France

¹³ Institute for Interdisciplinary Mountain Research, Technikerstrasse 21a, ICT Gebäude, 6020 Innsbruck, Austria

¹⁴ 8952 Irnding-Donnersbachtal, Raumberg 38, Austria

¹⁵ Experimental sPlant Ecology, Institute of Botany and Landscape Ecology, University of Greifswald, Soldmannstr. 15; Room 1.02, 17487 Greifswald, Germany

¹⁶ BioSense Institute, Dr Zorana Djindjica 1, Novi Sad, Serbia

¹⁷ Thoday Building, School of Environment, Natural Resources & Geography, Bangor University, Bangor LL57 2UW, UK

¹⁸ EnvixLab, Dipartimento di Bioscienze e Territorio, Università degli Studi del Molise, Via Duca degli Abruzzi s.n.c., 86039 Termoli, Italy

¹⁹ Dept. Zoology and Animal Ecology, Fac. of Environmental Sciences, SZIE University, 2100 Gödöllő, Páter K. 1., Hungary

²⁰ Helmholtz Centre for Environmental Research - UFZ, Department of Community Ecology, Theodor-Lieser-Straße 4, 06120 Halle, Germany

²¹ CREAF, Campus Universitat Autònoma de Barcelona, Edifici C, 08193 Bellaterra, Spain

²² ECOLAB, CNRS-UPS-INPT, ENSAT Avenue de l'Agrobiopole, BP, 32607, Auzeville-Tolosane, 31326 Castanet-Tolosan, France

²³ Centre for Biodiversity & Taxonomy, Department of Botany, University of Kashmir, Srinagar 190 006, Jammu, and Kashmir, India

²⁴ Instituut voor Natuur-en Bosonderzoek (INBO), Gaverstraat 4, 9500 Geraardsbergen, Belgium

²⁵ Forest Resources Department, University of Minnesota, 1530 Cleveland Ave. N, St. Paul, MN 55720, USA

²⁶ Dept. Ecología-Pontificia Universidad Católica de Chile & Instituto de Ecología y Biodiversidad, Alameda 340, Santiago, Chile

²⁷ KU Leuven, Division of Forest, Nature and Landscape, Celestijnenlaan 200E, 3001 Leuven, Belgium

²⁸ University of Liège, Plant and Microbial Ecology, Botany B22, Quartier Vallée 1, Chemin de la Vallée 4, 4000 Liège, Belgium

²⁹ Northwest German Forest Research Institute, Grätzelstr. 2, D-37079 Göttingen, Germany

³⁰ Ontario Forest Research Institute, 1235 Queen St. E., Sault Ste. Marie, Ontario P6A 2E5, Canada

³¹ University of Innsbruck, Institute of Ecology, Technikerstrasse 25, 60 Innsbruck, Austria

³² Department of Forest Sciences, PO Box 27, 00014, University of Helsinki, Finland

³³ Key Laboratory of Plant Resources and Biodiversity of Jiangxi Province, Jingdezhen University, 838 Cidu Avenue, Jingdezhen, Jiangxi 333000, China

³⁴ Martin Luther University Halle-Wittenberg, Institute of Biology/Geobotany and Botanical Garden, Am Kirchor 1, 06108 Halle (Saale), Germany

³⁵ Katedra Pedológie, Prírodovedecká fakulta UK, Mlynská dolina, Ilkovičova 6, 842 15 Bratislava 4, Slovakia

³⁶ Institute of Botany, Sternwartestr. 15A, 6020 Innsbruck, Austria

- ³⁷ Universidade Federal de Mato Grosso, Instituto de Biotecnologia, Departamento de Botânica, Av. Fernando, Corrêa da Costa, no 2367, Bairro Boa Esperança, CEP 78060-900 Cuiabá, MT, Brazil
- ³⁸ Uni Research Climate, Jahnebakken 5, 5007 Bergen, Norway
- ³⁹ University of Vermont, Rubenstein School of Environment and Natural Resources, Aiken Forestry Science Lab, 705 Spear Street, South Burlington, VT 05403, USA
- ⁴⁰ INRA Biogeco, Pierroton, 33612 Cestas, France
- ⁴¹ University of Freiburg, Geobotany, Schänzlestr. 1, 79104 Freiburg, Germany
- ⁴² Division of Agricultural Chemistry, Taiwan Agricultural Research Institute (TARI), Council of Agriculture, Executive Yuan, No. 189, Zhongzheng Rd., Wufeng Dist., Taichung City 41362, Taiwan
- ⁴³ Division of Silviculture, Soil Lab., Nan-Hai Rd. No. 53, Taipei, Taiwan
- ⁴⁴ Lehrstuhl für Bodenkunde, Agrar- und Umweltwiss. Fakultät, Justus-von-Liebig Weg 6, D-18059 Rostock, Germany
- ⁴⁵ WSL Institute for Snow and Avalanche Research SLF, Flüelastasse 11, 7260 Davos, Dorf, Switzerland
- ⁴⁶ CNRM, CNRS - Météo France, 42 av. G. Coriolis, 31057 Toulouse Cedex, France
- ⁴⁷ CEH, Bush Estate, Penicuik EH26 0QB, UK
- ⁴⁸ Department of Geosciences and Natural Resource Management, University of Copenhagen, Rolighedsvej 23, 1958 Frederiksberg, Denmark
- ⁴⁹ Helmholtz Centre for Environmental Research - UFZ, Department Computational Hydrosystems, Permoser Str. 15, 05318 Leipzig, Germany
- ⁵⁰ Centre for Ecology, Evolution and Environmental Changes (cE3c), Faculdade de Ciências, Universidade de Lisboa, 1749-016, Lisboa, Portugal
- ⁵¹ State Nature Reserve "Stolby", Kariymaya Str. 26a, Krasnoyarsk RU660006, Russia
- ⁵² Department of Geography, University of Bonn, Meckenheimer Allee 166, D-53115 Bonn, Germany
- ⁵³ Soil & Hydrology Research, AL-Quds University, P.O. Box 89, Bethlehem, Palestine
- ⁵⁴ Salah Al-Din st., East Jerusalem, P.O. Box: 67743, Israel
- ⁵⁵ Departamento de Química, Instituto Nacional de Investigaciones Nucleares, Carr. Mexico-Toluca, S/N, La Marquesa, Ocoyoacac, Estado de Mexico, Mexico
- ⁵⁶ Virginia Commonwealth University Rice Rivers Center, 3701 John Tyler Memorial Hwy, Charles City County, VA 23030, USA
- ⁵⁷ Research Centre in Systems Ecology and Sustainability, Faculty of Biology, University of Bucharest, Splaiul Independentei 91-95, 050095, District 5, Bucharest, Romania
- ⁵⁸ Dead Sea and Arava Science Center, P.O. Box 262, Mitzpe Ramon, Israel
- ⁵⁹ Chrono-Environnement, CNRS-Université de Bourgogne Franche-Comté & LTZER Zone Atelier Arc Jurassien, 16 route de Gray, 25030 Besançon Cedex, France
- ⁶⁰ Dept. Ecology, Inst. Biology, University of Veterinary Medicine, Rottenbiller u. 50, 1077 Budapest, Hungary
- ⁶¹ Centre d'Estudis Avançats de Blanes-CSIC, Ctra Accés Cala St. Francesc 14, 17300 Blanes, Spain
- ⁶² Université du Québec à Trois-Rivières, Trois-Rivières, Quebec, Canada
- ⁶³ Universidade Estadual de Maringá, Nupelia, Av. Colombo, 5790, 87020-900 Maringá, PR, Brazil
- ⁶⁴ Altai State University, Lenina Ave. 61, Barnaul RU-656049, Russia
- ⁶⁵ Tigirek State Reserve, Nikitina Str. 111-42, Barnaul RU-656043, Russia
- ⁶⁶ Ghent University, Forest & Nature Lab, Campus Gontrode, Geraardsbergsesteenweg 267, 9090 Melle-Gontrode, Belgium
- ⁶⁷ Universidade Federal de Mato Grosso do Sul, Centro de Ciências Biológicas e da Saúde, 79070-900 Campo Grande, MS, Brazil
- ⁶⁸ Herbario QCA, Departamento de Biología Pontificia Universidad Católica del Ecuador, Av. 12 de Octubre, entre Patria y Veintimilla, Apartado 17-01-2184, Quito, Ecuador
- ⁶⁹ Federal Research and Training Centre for Forests, Natural Hazards and Landscape (BFW), 1131 Wien, Seckendorff-Gudent-Weg 8, Austria
- ⁷⁰ Universidad Rey Juan Carlos Departamento de Biología y Geología, Física y Química Inorgánica Escuela Superior de Ciencias Experimentales y Tecnología C/ Tulipán s/n, Móstoles 28933, Spain
- ⁷¹ LIEC, CNRS-Université de Lorraine & LTZER Zone Atelier du Bassin de la Moselle, Campus Bridoux - Avenue du Général Delestraint, 57070 Metz, France
- ⁷² Department of Botany & Biodiversity Research, Division of Conservation Biology, Vegetation- and Landscape Ecology, University of Vienna, Austria
- ⁷³ Universität Innsbruck, Institut für Ökologie, Sternwartestr. 15, 6020 Innsbruck, Austria
- ⁷⁴ Swiss Federal Research Institute WSL, Zuercherstrasse 111, 8903 Birmensdorf, Switzerland
- ⁷⁵ Biodiversity Department - Consorcio para el Desarrollo Sostenible de la Ecorregión Andina (CONDESAN), Germán Alemán E12-123, Quito, Ecuador
- ⁷⁶ Palaeoecology & Landscape Ecology, Institute for Biodiversity & Ecosystem Dynamics (IBED), University of Amsterdam, Netherlands
- ⁷⁷ Universidade Federal de Mato Grosso, Faculdade de Agronomia, Medicina Veterinária e Zootecnia, Departamento de Solos e Engenharia Rural, Av. Fernando Corrêa, no 2367, Campus Universitário, Bairro Boa Esperança, CEP: 78060-900 Cuiabá, MT, Brazil
- ⁷⁸ Institute of Soil Research, University of Natural Resources and Life Sciences, Peter-Jordan-Str. 82, 1190 Vienna, Austria
- ⁷⁹ Galapagos National Park Directorate, Puerto Ayora, Santa Cruz, Island, Galapagos, Ecuador
- ⁸⁰ Department of Biology, California State University, 333 S. Twin Oaks Valley Road, San Marcos, CA 92096, USA
- ⁸¹ CATIE, Agroforesteria, DID. Cratago, Turrialba, Turrialba 30501, Costa Rica
- ⁸² The World Agroforestry Centre, Latin America Regional Office, Central America, CATIE 7170, Turrialba 30501, Cartago, Costa Rica
- ⁸³ Plant Conservation Unit, Department of Biological Sciences, University of Cape Town, Rondebosch, 7701 Cape Town, South Africa
- ⁸⁴ USFS International Institute of Tropical Forestry, 1201 Calle Ceiba, San Juan 00926, Puerto Rico
- ⁸⁵ Computational and Applied Vegetation Ecology Lab, Department of Applied Ecology and Environmental Biology, Ghent University, Coupure Links 653, 9000 Gent, Belgium
- ⁸⁶ German Centre for Integrative Biodiversity Research, (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany
- ⁸⁷ GLORIA-Coordination, Austrian Academy of Sciences (IGF), Austria
- ⁸⁸ University of Natural Resources and Life Sciences Vienna (ZgWN), Silbergasse 30/3, 1190 Vienna, Austria
- ⁸⁹ Université Nazi Boni, Institut du Développement Rural, Laboratoire d'étude et de Recherche sur la Fertilité du sol, BP 1091, Bobo-Dioulasso, Burkina Faso
- ⁹⁰ INTA, Casilla de Correo 332, CP 9400 Río Gallegos, Santa Cruz, Argentina
- ⁹¹ Animal Ecology I, Bayreuth Center for Ecology and Environmental Research (BayCEER), University of Bayreuth, 95440 Bayreuth, Germany
- ⁹² Charles Darwin Foundation, Puerto Ayora, Galápagos Islands, Ecuador
- ⁹³ CEFS, INRA, 24 Chemin de Borde Rouge, Auzeville, CS, 52627, 31326 Castanet-Tolosan Cedex, France
- ⁹⁴ Northwest German Forest Research Institute, Grätzelstrasse 2, 37079 Göttingen, Germany
- ⁹⁵ University of Landau, Fortstr. 7, 76829 Landau, Germany
- ⁹⁶ Forest Research Station, Field Science Center for Northern Biosphere, Hokkaido University, Kita-9, Nishi-9, Kita-ku, Sapporo 060-0809, Japan
- ⁹⁷ National Research and Development Agency, Forest Research and Management Organization, Forestry and Forest Products Research Institute, 1 Matsunosato, Tsukuba 305-8687, Japan
- ⁹⁸ Estación El Carmen, ICMYL, UNAM, km 9.5 Carretera Carmen-Puerto Real, Ciudad del Carmen, Campeche 24157, Mexico
- ⁹⁹ Universidad Autónoma de Ciudad Juárez (UACJ), Sede Cuauhtémoc-Programa de Geoinformática, Km. 3.5 Carretera Anáhuac, Municipio de Cuauhtémoc, Chihuahua CP 31600, Mexico
- ¹⁰⁰ United States Department of Agriculture Forest Service, 5523 Research Park, Suite 350, Baltimore, MD 21228, USA
- ¹⁰¹ Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse 111, 8903 Birmensdorf, Zürich, Switzerland
- ¹⁰² Institute of Biology, University of Latvia, Miera str. N3, Salaspils, LV -2169, Latvia
- ¹⁰³ Plant Department, Faculty of Science, University of Kisangani, People's Republic of Congo
- ¹⁰⁴ Centre Universitaire Polytechnique de Dédougou-UO I Pr Joseph KI-ZERBO, Laboratoire d'étude et de Recherche sur la fertilité du sol (UNB), 01, BP 7021, Ouagadougou, Burkina Faso
- ¹⁰⁵ Institute of Environmental Sciences, University of Nyíregyháza, Sóstói u. 31./B., 4400 Nyíregyháza, Hungary
- ¹⁰⁶ Institute of Forest Ecology, Slovak Academy of Sciences, L. Stura 2, 96053 Zvolen, Slovakia
- ¹⁰⁷ Institute of Ecology and Earth Sciences, University of Tartu, Vanemuise 46, 51014 Tartu, Estonia
- ¹⁰⁸ Ústav krajinnnej ekológie SAV, Pobočka Nitra, Akademická 2, P.O. Box 22, 949 10 Nitra, Slovakia
- ¹⁰⁹ Departamento de Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, Ctra. Utrera km. 1, 41013 Sevilla, Spain
- ¹¹⁰ Sorbonne Universités, UPMC Univ Paris 06, CNRS, INRA, IRD, Univ Paris Diderot Paris 07, UPEC, UMR 7618, Institute of Ecology and Environmental Sciences, Paris, France
- ¹¹¹ Centre Alpin de Phytogéographie, Fondation J.-M. Aubert, 1938 Champex-Lac, Switzerland
- ¹¹² Section de Biologie, Université de Genève, Case postale 60, 1292 Chambésy, Switzerland
- ¹¹³ College of Natural Sciences, Department of Environmental Sciences, University of Puerto Rico-Río Piedras, P.O. Box 70377, San Juan 00936-8377, Puerto Rico
- ¹¹⁴ Forest Plantation Programme, Forest Research Institute Malaysia (FRIM), 52109 Kepong, Selangor, Malaysia
- ¹¹⁵ Marine Biological Laboratory, 7 MBL Street, Woods Hole, MA 02543, USA

- ¹¹⁶ Centre Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK
- ¹¹⁷ Institute of Botany, Dukelská 135, CZ-37982 Trebon, Czech Republic
- ¹¹⁸ Unidad Académica Procesos Oceánicos y Costeros, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Ciudad Universitaria, 04510 Ciudad de México, Mexico
- ¹¹⁹ South African Environmental Observation Network, Arid Lands Node, Kimberley 8306, South Africa
- ¹²⁰ Cedar Point Biological Station, 100 Cedar Point Road, Ogallala, NE 69153, USA
- ¹²¹ Lammi Biological Station, Pääjärventie 320, 16900 Lammi, Finland
- ¹²² Rif Field Station, Aðalbraut 16, 675 Raufarhöfn, Iceland
- ¹²³ CONACYT - Estación el Carmen, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Carretera Carmen-Puerto Real km. 9.5, 24157 Ciudad del Carmen, Campeche, Mexico
- ¹²⁴ E-22700 Jaca, Huesca, Spain
- ¹²⁵ Universidade Estadual de Londrina, CCB, BAV, Caixa Postal 10.011, 86.057-970 Londrina, PR, Brazil
- ¹²⁶ VCU Department of Biology, 1000 West Cary St., Richmond, VA 23284, USA
- ¹²⁷ Instituto Pirenaico de Ecología (IPE-CSIC), Avda. Llano de la Victoria 16, Jaca 22700 (Huesca), Spain
- ¹²⁸ Programa de Investigación en Biodiversidad y Recursos Ecosistémicos, Universidad Nacional de Loja, Ciudadela Universitaria, sector La Argelia, EC110101 Loja, Ecuador.
- ¹²⁹ Department of Biological and Environmental Sciences, College of Arts and Sciences, Qatar University, P.O. Box 2713, Doha, Qatar
- ¹³⁰ Eurac, Institut für Alpine Umwelt, Drususallee 1, 39100 Bozen, Italy
- ¹³¹ Université Nazi Boni, Institut du développement rural, Laboratoire d'étude et de recherche sur la fertilité du sol, BP 1091, Bobo-Dioulasso, Burkina Faso
- ¹³² Nakagawa Experimental Forest, Field Science Center for Northern Biosphere, Hokkaido University, 483 Ootoinappu, Ootoinappu 2501, Japan,
- ¹³³ Department of Earth and Planetary Sciences, Johns Hopkins University, 3400 N. Charles St, Baltimore, MD 21218, USA
- ¹³⁴ Dept. Crop and Soil Science, Oregon State University, Corvallis, OR 97330, USA
- ¹³⁵ Soil Science and Geomorphology, Institute of Geography, University of Tübingen, Rümelinstrasse 19-23, 72070 Tübingen, Germany
- ¹³⁶ 475 Morse Hall, 8 College Road, Durham, NH 03824, USA
- ¹³⁷ Plant Ecology Group, University of Tübingen, Auf der Morgenstelle 5, 72076 Tübingen, Germany
- ¹³⁸ Department of Biological Environment, Akita Prefectural University, Shimoshinjo, Akita 010-0195, Japan
- ¹³⁹ National Parks and Wildlife Service, P.O. Box 2228, Jindabyne, NSW 2627, Australia
- ¹⁴⁰ Forest & Nature Lab, Department of Forest and Water Management, Ghent University, Geraardsbergsesteenweg 267, 9090 Gontrode, Belgium
- ¹⁴¹ Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN 55108, USA
- ¹⁴² IFEVA, Catedra de Ecología, Facultad de Agronomía, UBA, Av. San Martin 4453, 1417 CABA, Argentina
- ¹⁴³ SUNY-ESF, Marshall Hall, 1 Forestry Drive, Syracuse, NY 13210, USA
- ¹⁴⁴ Plant Ecology Group, University of Tübingen, Tübingen, Germany
- ¹⁴⁵ USDA Forest Service Northern Research Station, 271 Mast Rd, Durham, NH 03824, United States
- ¹⁴⁶ Univeristà degli studi di Sassari, Dipartimento di Scienze per la Natura e il Territorio, via Enrico de Nicola 9, 07100 Sassari, Italy
- ¹⁴⁷ Majella Seed Bank, Majella National Park, Colle Madonna, 66010 Lama dei Peligni, Italy
- ¹⁴⁸ Ecological Plant Geography, Faculty of Geography, University of Marburg, Deutschhausstraße 10, DE-35032 Marburg, Germany
- ¹⁴⁹ ECOBIO CNRS-Université de Rennes 1 & LTSEZ Zone Atelier Antarctique et Subantarctique, Station Biologique, 35380 Paimpont, France
- ¹⁵⁰ Tel Aviv University, School of Plant Sciences and Food Security, Tel Aviv, Israel
- ¹⁵¹ Forest Research Institute - Bulgarian Academy of Sciences, 132 "St. Kl. Ohridski" blvd., 1756 Sofia, Bulgaria
- ¹⁵² K.-C.-Irving Chair in Environmental Sciences and Sustainable Development, Université de Moncton, Moncton, NB E1A 3E9, Canada
- ¹⁵³ 1910 University drive, Boise, ID 83703, United States
- ¹⁵⁴ LRGP, CNRS-Université de Lorraine & LTSEZ Zone Atelier du Bassin de la Moselle, 1 rue Grandville, BP, 20451, 54001 Nancy, cedex, France
- ¹⁵⁵ Isotope Bioscience Laboratory - ISOFYS, Department of Applied Analytical and Physical Chemistry, Ghent University, Coupure Links 653, 9000 Gent, Belgium
- ¹⁵⁶ Programa de Biodiversidad y Servicios Ecosistémicos, Universidad Nacional de Loja, Ciudadela Universitaria, sector La Argelia, EC110101 Loja, Ecuador
- ¹⁵⁷ Swiss Federal Research Institute WSL, Forest Resources and Management, Birmensdorf CH-8903, Switzerland
- ¹⁵⁸ Department of Biological Sciences, Montana Tech of the University of Montana, Butte, MT 59701, USA
- ¹⁵⁹ Swedish University of Agricultural Sciences, Department of Crop Production Ecology, PO Box 7043, SE-75007 Uppsala, Sweden
- ¹⁶⁰ Earth Systems Research Center, 475 Morse Hall, 8 College Road, Durham, NH 03824, USA
- ¹⁶¹ Reichergasse 48, 3411 Klosterneuburg-Weidling, Austria
- ¹⁶² LIEC, CNRS-Université de Lorraine & LTSEZ Zone Atelier du Bassin de la Moselle, Campus Bridoux - Avenue du général Delestraint, France
- ¹⁶³ TERN, College of Science & Engineering, James Cook University, Cairns, Australia
- ¹⁶⁴ Meynertgasse 32, 3400 Klosterneuburg, Austria
- ¹⁶⁵ Czech Geological Survey, Geologická 6, Prague, 5 152 00, Czech Republic
- ¹⁶⁶ Department of Physical Geography and Geocology, Faculty of Science, Charles University, Albertov 6, 128 43 Prague, Czech Republic
- ¹⁶⁷ Department of Life Health and Environmental Sciences, University of L'Aquila, Via Vetoio, loc. Coppito, 67100 L'Aquila, Italy
- ¹⁶⁸ Forest Tree Breeding Center, Forestry and Forest Products Research Institute, Hitachi, Ibaraki 319-1301, Japan
- ¹⁶⁹ Department of Taxonomy and Ecology, Faculty of Biology and Geology, "A. Borza" Botanical Garden, Babeş-Bolyai University, 42 Republicii Street, 400015 Cluj-Napoca, Romania
- ¹⁷⁰ Forestry and Forest Products Research Institute (FFPRI), 68 Nagaikyutaroh, Momoyama, Fushimi, Kyoto 612-0855, Japan
- ¹⁷¹ Forest Research, Alice Holt Lodge, Farnham, Surrey GU10 4LH, United Kingdom
- ¹⁷² Field Science Education and Research Center, Kyoto University, Kyoto 606-8502, Japan
- ¹⁷³ Canada Research Chair in Polar and Boreal Ecology, Department of Biology, Université de Moncton, Moncton, NB E1A 3E9, Canada
- ¹⁷⁴ Yugra State University, 628508, Stroiteley Street, 2, Shapsha village, Khanty-Mansiyskiy rayon, Tyumen Region, Russia
- ¹⁷⁵ Institute of Landscape Ecology, University of Münster, Heisenbergstraße 2, 48149 Münster, Germany
- ¹⁷⁶ Bryn Mawr College, 101 N. Merion Ave., Bryn Mawr, PA 19010, USA
- ¹⁷⁷ Institute of Earth Surface Dynamics, University of Lausanne, Géopolis, 1015 Lausanne, Switzerland
- ¹⁷⁸ "A. Borza" Botanical Garden, Babeş-Bolyai University, 42 Republicii Street, 400015 Cluj-Napoca, Romania
- ¹⁷⁹ Research station and Museum of Tatra National Park, Tatranska Lomnica, 059 60 Vysoké Tatry, Slovakia
- ¹⁸⁰ School of Life and Environmental Sciences, Centre for Integrative Ecology, Deakin University, Victoria 3216, Australia
- ¹⁸¹ Department of River Ecology and Conservation, Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen, Germany
- ¹⁸² Faculty of Biology, University of Duisburg-Essen, Essen, Germany
- ¹⁸³ Department of Forest Resources, University of Minnesota, 1530 Cleveland Ave. N, St. Paul, MN 55108, USA
- ¹⁸⁴ Hawkebury Institute for the Environment, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751, Australia
- ¹⁸⁵ Institute of Botany, The Czech Academy of Sciences, Zámek 1, 25243 Pílhonice, Czech Republic
- ¹⁸⁶ LEHNA, CNRS-Université Claude Bernard 1 & LTSEZ Zone Atelier Bassin du Rhône, 43 Boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France
- ¹⁸⁷ Earth and Life Institute, Université catholique de Louvain, Croix du Sud 2 - Box L7.05.09, 1348 Louvain-la-Neuve, Belgium
- ¹⁸⁸ Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Kreuzeckbahnstr. 19, 82467 Garmisch-Partenkirchen, Germany
- ¹⁸⁹ Center for Ecosystem Science and Society, Northern Arizona University, P.O. Box 5620, Flagstaff, AZ 86011, USA
- ¹⁹⁰ Arctic Station, University of Copenhagen, 3953 Qeqertarsuaq, Greenland
- ¹⁹¹ Leibniz Institute of Freshwater Ecology and Inland Fisheries Berlin, Müggelseedamm 301, 12587 Berlin, Germany
- ¹⁹² Ústav krajinnnej ekológie SAV, Štefánikova 3, 814 99 Bratislava, Slovakia
- ¹⁹³ Experimental Plant Ecology, Institute of Botany and Landscape Ecology, University of Greifswald, Soldmannstr. 15, 17487 Greifswald, Germany

- ¹⁹⁴ Bio-Clim-Land Centre, Biological Institute, Tomsk State University, Tomsk, Russia
- ¹⁹⁵ NUPEM, Federal University of Rio de Janeiro (UFRJ), Av. São José do Barreto, 764, B. São José do Barreto, Postal Code 27965-045 Macaé, RJ, Brazil
- ¹⁹⁶ Museo Nacional de Historia Natural, Herbario Nacional de Bolivia, Campus Universitario UMSA, Cota Cota calle 27, La Paz, Bolivia
- ¹⁹⁷ Department of Forestry and Natural Resources, Chauras Campus, H.N.B. Garhwal University (A central University), Post Office: Kilkleshwar, Kirtinagar, Tehri Garhwal, Uttarakhand 249161, India
- ¹⁹⁸ Wrigley Global Institute of Sustainability, Arizona State University, PO Box 875402 (800 S. Cady Mall), Tempe, AZ 85287-5402, USA
- ¹⁹⁹ The University of Tokyo Chichibu Forest, The University of Tokyo, 1-1-49 Hinoda-machi, Chichibu, Saitama 368-0034, Japan
- ²⁰⁰ Florida International University Biology Department, OE 00148, 11200 SW 8th Street, Miami, FL 33199, USA
- ²⁰¹ Observatoire des Sciences de l'Univers en région Centre – OSUC, Service d'Observation Tourbières, Campus Géosciences, 1A rue de la Férolerie, 45071 Orléans Cedex, France
- ²⁰² Geoecology, Department of Geography and Regional Research, University of Vienna, Althanstraße 14, AT-1090 Vienna, Austria
- ²⁰³ Euro-Mediterranean Center on Climate Change, Impacts on Agriculture, Forests and Natural Ecosystems (IAFES) Division, Sassari, Italy
- ²⁰⁴ Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, 221 Burwood Hwy, Burwood, VIC 3125, Australia
- ²⁰⁵ Department of Environmental Science and Assessment, Swedish University of Agricultural Sciences SLU, P.O. Box 7050, SE-750 07 Uppsala, Sweden
- ²⁰⁶ University of Applied Sciences Trier, Umwelt-Campus Birkenfeld, Postbox 1380, 57761 Birkenfeld, Germany
- ²⁰⁷ DST/NRF Centre of Excellence at the Percy FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch 7701, South Africa
- ²⁰⁸ LIEC, CNRS-Université de Lorraine & LTSEZ Zone Atelier du Bassin de la Moselle, BP 70239, Bd des Aiguillettes, 54506 Vandœuvre-les-Nancy, France
- ²⁰⁹ South China Botanical Garden, Chinese Academy of Sciences, Xingke Road #723, Tianhe District, Guangzhou, Guangdong 510650, China
- ²¹⁰ Ecohydrology Research Institute, The University of Tokyo Forests, 11-44 Goizuka, Seto, Nagoya, Aichi 489-0031, Japan
- ²¹¹ Shiiba Research Forest, Kyushu University, 949 Ohkawachi, Shiiba Village, Prefecture Miyazaki 883-0402, Japan
- ²¹² 1278-294 Sugadaira-kogen, Ueda 386-2204, Japan
- ²¹³ Tomakomai Experimental Forest, Hokkaido University, Takaoka, Tomakomai, Hokkaido 053-0035, Japan
- ²¹⁴ Laboratório de Ecologia de Comunidades, Instituto de Ciências Biológicas, Universidade Federal do Pará, Rua Augusto Correia, No 1, P.O. Box: 479, Zip Code 66075-110 Bairro Guamá, Belém, Pará, Brazil
- ²¹⁵ ECOLAB, INPT-ENSAT, Avenue de l'Agrobiopole, BP 32607, 31326 Castanet Tolosan Cedex, France
- ²¹⁶ Univ. Grenoble Alpes, Irstea, EMGR, LTSEZ Zone Atelier Alpes, F-38000 Grenoble, France
- ²¹⁷ Biological Station Lake Neusiedl, 142 Illmitz, Seewogelände 1, Austria
- ²¹⁸ Kasuya Research Forest, Kyushu University, 394 Tsubakuro, Sasaguri, Fukuoka 811-2415, Japan
- ²¹⁹ Institute of Biological Research, Department of Taxonomy and Ecology, National Institute of Research and Development for Biological Sciences, 400015 Cluj-Napoca, Romania
- ²²⁰ Regional Environmental Protection Agency - Aosta, Valley, Loc. Grande Charrière, 44, Saint-Christophe 11020 - I, Italy
- ²²¹ Universidade Federal de Mato Grosso, Instituto de Biotecnologia, Dourados PPG Ecologia e Conservação da Biodiversidade, Av. Fernando Corrêa da Costa, no 2367, Bairro Boa Esperança, CEP 78060-900 Cuiabá, MT, Brazil
- ²²² Mazingira Centre, International Livestock Research Institute (ILRI), P.O. Box 30709, 00100 Nairobi, Kenya
- ²²³ CEB-CNRS & LTSEZ Zone Atelier Plaine et Val de Sèvre, 79360 Beauvoir sur Niort, France
- ²²⁴ Dept. Ecologia, Inst. Biologia, CCS, Bloco A, Sala A0-008, Ilha do Fundão, Rio de Janeiro, RJ, 21941-590, Brazil
- ²²⁵ 88th, Xuefu, Road, Kunming, Yunnan Province 650223, China
- ²²⁶ Erguna Forest-Steppe Ecotone Research Station, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110164, China
- ²²⁷ Natural Resources and the Environment, University of New Hampshire, Durham, NH 03824, USA
- ²²⁸ Fujian Agricultural & Forestry University, No. 15 Shangxiadian Road, Fuzhou 350002, China
- ²²⁹ Ashoro Research Forest, Kyushu University, 1-85 Kita 5, Ashoro, Ashoro-gun, Hokkaido 089-3705, Japan
- ²³⁰ Khibiny Research and Educational station of the Faculty of Geography, Lomonosov Moscow State University, ul. Zheleznodorozhnaya 10, Kirovsk 184250, Murmansk region, Russia
- ²³¹ State Nature Reserve "Olekminsky", Filatova Str. 6, Olekminsk, Yakutia Ru-678100, Russia
- ²³² Department of Soil Science and Water Management, Szent István University of Budapest, H-1118, Budapest, Villányi út. 29-43, Hungary
- ²³³ Current address: Institute of Agricultural Sciences, ETH Zurich, Universitätsstr. 2, 8092 Zurich, Switzerland

^a Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland

^b Department of Geosciences and Natural Resource Management, University of Copenhagen, Rolighedsvej 23, 1958 Frederiksberg, Denmark


^c Department of Forest Sciences, University of Helsinki, Latokartanonkaari 7, 00014 Helsinki, Finland

^d Forest & Nature Lab, Department of Forest and Water Management, Ghent University, Geraardsbergsesteenweg 267, 9090 Gontrode, Belgium

HIGHLIGHTS

- Litter quality is the key driver of initial litter decomposition at the global and regional scale.
- MAT has a low explanatory power on initial litter decomposition and is litter specific.
- MAP significantly affected litter decomposition but has low explanatory power.
- When data were aggregated at the biome scale, climate played a significant role on decomposition.
- The TeaComposition initiative is a low-cost standardized metric on litter decomposition.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 31 August 2017

Received in revised form 20 December 2017

Accepted 2 January 2018

Available online xxxx

ABSTRACT

Through litter decomposition enormous amount of carbon is emitted to the atmosphere. Numerous large-scale decomposition experiments have been conducted focusing on this fundamental soil process in order to understand the controls on the terrestrial carbon transfer to the atmosphere. However, previous studies were mostly based on site-specific litter and methodologies, adding major uncertainty to syntheses, comparisons and meta-analyses across different experiments and sites. In the TeaComposition initiative, the potential litter decomposition is investigated by using standardized substrates (Rooibos and Green tea) for comparison of litter mass loss in

Keywords:

Tea bag
Green tea
Rooibos tea
Carbon turnover
TeaComposition initiative

336 sites (ranging from -9 to $+26$ °C MAT and from 60 to 3113 mm MAP) across different ecosystems. In this study we tested the effect of climate (temperature and moisture), litter type and land-use on early stage decomposition (3 months) across nine biomes. We show that litter quality was the predominant controlling factor in early stage litter decomposition, which explained about 65% of the variability in litter decomposition at a global scale. The effect of climate, on the other hand, was not litter specific and explained $<0.5\%$ of the variation for Green tea and 5% for Rooibos tea, and was of significance only under unfavorable decomposition conditions (i.e. xeric versus mesic environments). When the data were aggregated at the biome scale, climate played a significant role on decomposition of both litter types (explaining 64% of the variation for Green tea and 72% for Rooibos tea). No significant effect of land-use on early stage litter decomposition was noted within the temperate biome. Our results indicate that multiple drivers are affecting early stage litter mass loss with litter quality being dominant. In order to be able to quantify the relative importance of the different drivers over time, long-term studies combined with experimental trials are needed.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

Through litter decomposition $>50\%$ of net primary production is returned to the soil (Wardle et al., 2004) and $60 \text{ Pg C year}^{-1}$ is emitted to the atmosphere (Houghton, 2007). Depending on the type of ecosystem, the quantity of soil organic carbon (SOC) in the top 1-m depth range from 30 tons/ha in arid climates to 800 tons/ha in organic soils in cold regions, with a predominant range from 50 to 150 tons/ha (Lal, 2004). The amount of SOC is determined by the balance of carbon inputs from primary production and losses through the decomposition of organic matter over time (Olson, 1963). However, there is a large degree of variability in this balance and more research is needed for a better mechanistic understanding of decomposition processes at various scales and for a more accurate estimation of present and future global carbon budgets (Aerts, 2006).

Decomposition of plant litter may be divided into at least two stages (e.g. Berg and McClaugherty, 2008). The early stage of decomposition (ca. 0 to 40% mass loss) is characterized by leaching of soluble compounds and by decomposition of solubles and non-lignified cellulose and hemicellulose (Couteaux et al., 1995; Heim and Frey, 2004). The late stage (ca. 40–100% mass loss) encompasses the degradation of lignified tissue. In general, microbial decomposition of organic substrates is controlled by both biotic factors (substrate quality and microbial community composition) and abiotic factors (temperature and moisture; Gavazov, 2010). Research to understand the impact of global changes such as climate on decomposition processes has typically been conducted at individual sites and/or through cross-site observations and experiments (e.g. Emmett et al., 2004; Heim and Frey, 2004; García Palacios et al., 2013). This has sometimes lead to controversial conclusions since the observed decomposition may be dependent on local litter quality used in the study and the factors controlling decomposition may be influenced by the methodologies and experimental designs applied. Consequently, comparisons across observations and common conclusions may be hampered. For example, early stage decomposition (mainly microbial) has been reported to be primarily controlled by climate and major nutrients in pine needle litter (Berg and McClaugherty, 2008), by microbial and nematode communities in pine needle litter (García Palacios et al., 2016), by litter content of water soluble substances (Heim and Frey, 2004) and by soil temperature and soil pH for a maize straw-soil mixture (Djukic et al., 2012). At regional and global scales, litter decomposition has been reported to be controlled by climate and litter quality (explaining about 60–70% of litter decomposition rates; Parton et al., 2007) and by soil meso- and micro-fauna communities (explaining about 7%; Wall et al., 2008). However, at the biome scale the metadata-analysis by García Palacios et al. (2013) showed that the variables controlling decomposition vary with decomposition in cold and dry biomes being mostly controlled by climatic conditions while soil fauna seemed to have a more defining role in warm and wet biomes. Moreover,

Bradford et al., (2014), showed that climate has a main control on decomposition only when local-scale variation is aggregated into mean values. In order to pinpoint the specific drivers of litter decomposition across various litter types with different decomposition rates and across multiple sites, standardized studies across sites and regions are needed (Wickings et al., 2012; Handa et al., 2014; Parsons et al., 2014).

Decomposition studies across multiple sites using standardized methods already exist within observational networks or experimental studies such as GLIDE (Global Litter Invertebrate Decomposition Experiment – Wall et al., 2008), LIDET (Long-term Intersite Decomposition Experiment Team – Adair et al., 2008), CIDET (Canadian Intersite Decomposition Experiment – Trofymow and CIDET Working Group, 1998), DIRT (Detrital Input and Removal Experiment – Nadelhoffer, 2004), BioCycle (Biodiversity and biogeochemical cycles: a search for mechanisms across ecosystems – Makkonen et al., 2012), DECO (European Decomposition project – Johansson et al., 1995), CANIF (Carbon and Nitrogen Cycling in Forest Ecosystems project – Persson et al., 2000), MICS (Decomposition of organic matter in terrestrial ecosystems: microbial communities in litter and soil – Cotrufo et al., 2000), VULCAN (Vulnerability assessment of shrubland ecosystems in Europe under climatic changes – Emmett et al., 2004) and VAMOS (Variation of soil organic matter reservoir – Cotrufo et al., 2000). Results from these have been used by predictive models such as Yasso07 (Tuomi et al., 2009) and in meta-analyses such as the ART-DECO project (Cornwell et al., 2008). These studies have all provided important information on the decomposition of litter, but have been limited to specific biomes or ecosystem types or have used site specific litter.

Therefore, despite the many efforts, a general understanding of the litter decomposition process and its driving factors is hampered by (1) use of site- or network/project-specific litters and methodologies (e.g. different study lengths, litter bag mesh sizes, incubation depths, litter type and litter mixes; García Palacios et al., 2013), and (2) the low number of global studies that go across all biomes (Bradford et al., 2016). This study presents results from the TeaComposition initiative which uses standard litters (tea bags – Keuskamp et al., 2013) and a common protocol allowing global and long-term application to overcome these limitations by providing standardized litter decomposition measurements across broad spatial scales. The study presents early stage litter mass loss across nine biomes with the aim to determine and compare globally the main drivers of decomposition at present climatic conditions. The early stage decomposition is generally expected to show greater mass loss rates and a dynamic response of mass loss to controlling factors (e.g. Heim and Frey, 2004; Pérez-Suárez et al., 2012). Therefore the specific objectives of the study were to estimate the variation in early stage mass loss of two litter types worldwide, to explore the linkage of early stage litter mass loss with key drivers (climate, litter type, land-use), and to explore whether the relative importance of the drivers differ between the litter types. Our

research questions are (1) does early stage litter mass losses of Green tea and Rooibos tea vary at the global scale due to the different litter qualities (Didion et al., 2016; Keuskamp et al., 2013), (2) are abiotic drivers controlling the initial stage of mass loss (Bradford et al., 2016) with temperature being the main regulating factor in the cold biomes and precipitation in the warmer biomes (Adair et al., 2008), and (3) does early stage litter mass losses vary between land-use types due to changes in the microclimates (Fig. 1).

2. Material and methods

2.1. Background of the TeaComposition initiative

The TeaComposition initiative was started in summer 2016. The main objective is to investigate long-term litter decomposition and its key drivers at present as well as under different future climate scenarios using a common protocol and standard litter (tea) across nine terrestrial biomes. It is one of the first comprehensive global studies on litter decomposition focusing on the litter decomposition in the topsoil and the degradation of the main litter components (lignin, cellulose and hemicellulose) to carbon dioxide and soluble or leachable compounds. As a collaborative network the TeaComposition initiative has involved a large number of international research projects and networks with observational or experimental approaches, which are relevant for increasing our mechanistic understanding of decomposition processes as well as for improving the predictive power of process-based models.

2.2. Study sites

The TeaComposition initiative comprises 570 sites across nine terrestrial biomes (Fig. 2). Here “biome” is defined as a region with specific macroclimate and its classification was done

according to Walter and Breckle (1999). In this study, data from 336 sites were used for analyses. Some of the sites included manipulation experiments (e.g. including treatment plots such as fertilizer addition or climate manipulation) in which case only the tea bags from the untreated control plots were used in the analyses. Sub-sites with different conditions (e.g. tree species diversity experiments or altitudinal gradients) were considered as single sites.

Overall, the sites represented all terrestrial biomes (Table 1) and each site provided information on location (i.e. coordinates), climate (averaged monthly or daily temperature (MAT) and cumulative precipitation (MAP)), vegetation type, and specific land-use (Table S2). Climate data were measured at the site or taken from nearby weather stations. In cases where no climate data were provided, data were extracted from worldclim (Fick and Hijmans, 2017). The mean annual air temperature (MAT) in our dataset ranges from -9 to $+26$ °C and the mean annual precipitation (MAP) from 60 to 3113 mm (Table 1; Site specific data can be found in the Table S2). Since sites were assigned to different land-use categories from different classification schemes, we reclassified them into five broader classes: arable, forest, grassland, shrubland and wetland based on the site description.

2.3. Method and study design

The TeaComposition initiative uses tea bags as a standardized metric for decomposition as proposed by Keuskamp et al. (2013), and applies a standardized protocol adapted to match global and long-term applications. The standardized protocol ensures: (i) use of the same batch of tea bags assuring the same substrate quality for all sites, (ii) harmonized start of the decomposition at the same season at the year for northern and southern hemisphere (i.e. start in summer; June–August in northern hemisphere and December–February in southern hemisphere), (iii) comparable




Fig. 1. Conceptual depiction of the main research questions. The temperature dependency across the temperature range (figure b) is arbitrary.

Global litter decomposition study - TeaComposition sites 2017




Fig. 2. Map showing the location of the 570 study sites involved in the TeaComposition initiative so far. Data from the sites with the red circles have been used in the present study. Data from Qatar come from [Alsafran et al., 2017](#). See [Tables 1](#) and [S2](#) for more detailed information. Classification of the biomes was according [Walter and Breckle \(1999\)](#). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

incubation depth at the upper 5 cm of the soil relevant for litter decomposition, and (iv) standardized and comparable incubation times covering both short and long term dynamics with incubation times extending to three years (sampling points after 3, 12, 24, and 36 months).

Two types of tea material with distinct qualities are being used; the Green tea viz. green leaves (*Camellia sinensis*; EAN no.: 8 722700 055525) with high cellulose content and expected fast decomposition, and rooibos tea (*Aspalanthus linearis*; EAN no.: 8 722700 188438) with high lignin content and expected slow decomposition ([Keuskamp et al., 2013](#)). The bag material is made of

woven nylon and has a mesh size of 0.25 mm allowing access of microfauna ([Bradford et al., 2002](#)) in addition to microbes and very fine roots. Before the start of the incubation all tea bags were oven-dried at 70 °C for 48 h and the initial weight was recorded (overall mean = 1.81 g, s.d. = 0.10). Each bag was identified with a unique number and was buried in the upper 5 cm of the top soil layer during summer seasons in both the northern and southern hemisphere. At least two homogenous areas (plots) were selected (at least 1 m apart) at each site. Two replicates of the two litter qualities (Green tea and Rooibos tea) were installed in each of the two blocks, resulting in minimum 4, maximum 250,

Table 1
Summarized general characteristics of the study sites used for the analysis within the TeaComposition initiative. Note: Detailed table on the single site characteristics can be found in the Supplementary material.

Biomes	Number of sites	Land use	Climate data (MAT / MAP)*
Arctic climate	4	Grassland	-9 to 5 / 237 to 709
Boreal climate	17	Boreal Forest, Shrubland, Grassland, Bog, Ecotone	-3 to 6 / 293 to 1015
Temperate climate	250	Agriculture, Forest, Shrubland, Grassland (Meadows), Wetland, Ecotone, alpine Grassland	-7 to 14 / 265 to 2140
Warm-temperate climate	13	Forest, Shrubland, Grassland, Wetland	6 to 21 / 955 to 3072
Arid-temperate climate	9	Desert, Shrubland, Grassland steppe, Ecotone	6 to 21 / 174 to 528
Mediterranean climate	13	Agriculture, Forest, Shrubland, Grassland, Wetland, Lake, Subalpine / Alpine Grassland	7 to 25 / 569 to 1627
Subtropical arid climate	15	Forest, Grassland, Wetland	15 to 24 / 60 to 412
Equatorial humid climate	6	Agriculture, Forest, Wetland (Mangrove, Freshwater Swamp), Ecotone	22 to 26 / 1298 to 3113
Semi-arid tropical climate	9	Agriculture, Forest, Shrubland, Grassland (Savanna), Wetland	11 to 26 / 636 to 1268

* MAT = Mean annual temperature; MAP = Mean annual precipitation.

and in average 8.33 bags of each tea type per site and sampling time. Tea bags were collected at all sites after a field incubation period of three months. The tea bags were cleaned from soil and roots, oven dried (70 °C for 48 h), and the weight of the remaining tea (without bag) was recorded. Instead of weighing incubated tea bags (as often damaged, tag dissolved or rope missing) an averaged bag weight (40 empty tea bags; 0.248 g per bag) was used to estimate the amount of the tea before the incubation. If the collected tea bags were visibly contaminated with soil, ash content (refers to the mineral residue after removal of organic matter by ignition) was determined by heating in a muffle oven at 500 °C for 16 h, in order to correct for the mineral part (Soil Survey Staff, 2004).

2.4. Data analyses

Because not all tea bags were incubated for exactly three months (overall mean = 92 days, s.d. = 13.2) we linearly standardized all mass loss data to a fixed period of 90 days prior to data analyses. As such, the reported mass loss data therefore represent a rate of mass loss over 90 days.

2.4.1. Differences in tea mass loss across biomes and between tea types

We quantified differences in remaining litter mass between biomes using linear mixed models with biome and tea type as fixed factors and site as a random factor accounting for the dependence in observations within site. Residual plots were visually inspected for deviations from model assumptions. If the interaction between biome and tea type was significant, multiple comparisons between biomes within each tea type were tested applying post hoc contrasts with P-values adjusted for multiplicity with the single-step method (Hothorn et al., 2008).

To quantify the different sources of variation in our data we used a linear mixed effect model with a nested structure (sites nested within biome). Biome and site were set as random factors and tea type as a fixed factor. We then ran separate analyses for each tea type to investigate whether biome, site and individual tea bags accounted differently for the variation for each tea type.

2.4.2. Effects of climate on the initial litter mass loss

To investigate the effects of climatic variables on remaining tea mass after three months of field incubation we applied linear mixed models with local climate as fixed factors and site as random factor. We used local climate data (average monthly air temperature and total precipitation) measured at nearby weather stations during the period of incubation when data were available ($n = 124$; Fig. 4; Table 2). For sites with no local climate data, we imputed the monthly averages of temperature and the total precipitation for the corresponding

measurement period from Worldclim (Fick and Hijmans, 2017). Whereas local climate represent the weather conditions measured at the sites during the incubation period, Worldclim represents the average climate for the period 1970–2000. We assessed the congruency between the two types of climate data by also running models including only the sites where both types of data were available. The results were qualitatively similar to the model including all sites. Moreover, local and Worldclim climate data were highly correlated (precipitation: $r = 0.83$; $P < .01$; temperature: $r = 0.87$, $P < .01$, Pearson's product moment correlation).

We modeled the remaining mass as a function of tea type, temperature and precipitation. Differences between litter types were tested by including interaction terms for tea type with both climatic variables. We used backward selection for model simplification until only significant terms remained in the final model. When a significant interaction with tea type was found, we used post hoc contrasts to test for significant relationships between the climatic variable and each tea type (i.e. test for slope different from 0); P-values were adjusted for multiplicity using a single-step method based on the joint normal distribution. Goodness of fit for these models were calculated based on marginal and conditional R^2 (Nakagawa and Schielzeth, 2013). Because climatic effects on decomposition can depend on the spatial scale of the observation (Bradford et al., 2014) we conducted a separate analysis, using the average remaining mass, temperature and precipitation, aggregated at the biome level. We tested for effects of climate factors using simple linear models, with temperature, precipitation and their interaction as independent variables. Significant interactions were further tested as described above.

2.4.3. Effects of land-use on the initial litter mass loss

We tested for differences in remaining tea mass between land-use types only for the temperate biome as this was the




Fig. 3. Percentage remaining mass for Green and Rooibos teas across climatic biomes. The difference between Tea types was significant ($F = 9802$; $P < .01$). Red and green circles show the mean and the bars are the standard errors based on the total number of observations. Letters show pairwise comparisons within each tea type: lowercase for rooibos and uppercase for green. Numbers in parentheses are the total number of tea bags for each biome. Biomes are ordered by increasing mean annual precipitation. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Table 2

Effects of climatic factors on the site level remaining mass of the two tea types (statistics relates to Fig. 4). Estimates obtained from mixed effect model with site as a random factor. R^2 marginal: 0.74; R^2 conditional = 0.88.

	Est.(SE) ^a	t	P
Green tea	45.81(1.79)	25.62	<.01
Rooibos tea	79.57(1.80)	44.31	<.01
PREC	−8.87(2.68)	−3.32	<.01
Green tea × TEMP	0.14(0.17)	0.88	.38
Rooibos tea × TEMP	−0.12(0.17)	−0.74	.82

^a Models were fitted using precipitation/1000 to avoid very small estimates. Est. = estimates, SE = standard error.

only biome with enough sites of the different land-use categories. We used a mixed model including land-use, tea type and their interaction as fixed factors and site as random factor. Separate models were used for each tea type to further explore differences. If the interaction between land-use type and tea type was significant, multiple comparisons among land-use types within each tea type were tested using post hoc contrasts with P-values adjusted for multiplicity with the single-step method.

All statistical analyses were conducted with R (version 3.1.2; R core team 2014). The level for detecting statistical differences was set at $P < .05$. The lme4 package (Bates et al., 2015) was used for fitting the mixed models and the multcomp package (Hothorn et al., 2008) was used for multiple comparisons. The percentage of variance explained by the fixed and the different random components was calculated using the “variancePartition” library in R (Hoffman and Schadt, 2016).

3. Results

3.1. Relative importance of litter quality on mass loss across biomes

Across all biomes, tea mass remaining after three months of field incubation (Fig. 3) was higher for Rooibos tea (78%, SD = 10.31) than for Green tea (38%, SD = 15.86). Overall, similar mass loss patterns were recorded for both tea types across biomes with tendencies or significantly higher mass loss at warm and humid climates compared to the dry and/or cold biomes. However, there was a significant interaction between biome and tea type ($F = 84$; $P < .01$) indicating that some differences between biomes depend on tea type. For Rooibos tea, significantly lower remaining mass was found at sites in equatorial-humid climate. For Green tea, we found the highest remaining mass at the sites from the arid-subtropical and Mediterranean climates, which were significantly different from the sites found in cooler and more humid biomes (Fig. 3).

The analysis of data variation showed that 65% of the variation in the remaining litter mass was related to tea type while 13% was related to biome (Fig. 3). The variation was 11% within biomes and 11% within sites.

3.2. Effects of climate on the initial litter mass loss

Our final model showed that climatic variables had different effects on early stage decomposition. Remaining mass loss decreased with increasing precipitation. This pattern was similar for both tea types as revealed by the not significant interaction between tea type and precipitation ($F = 0.01$, $P = .96$). We also found a significant interaction between tea type and temperature ($F = 64$, $P < .01$) indicating that the response of mass loss to temperature depends on tea type, i.e. litter quality. However, the analyses using post hoc contrasts showed that temperature did not have any significant effect on any of the tea types (Table 2; Fig. 4).

In contrast, the biome-scale analyses focusing on the mean values for the given biome revealed some variation in remaining litter mass loss from low (equatorial humid climate) to high (arid subtropical and Mediterranean climates) mass losses (Fig. 5a). In the linear models, we found a non-significant interaction between tea type and MAP ($F = 0.20$, $P = .66$); and between tea types and MAT ($F = 0.39$, $P = .54$). Whereas MAT had no effect ($F = 0.64$, $P = .43$), remaining mass decreased with increasing MAP for both tea types (Table 3).

3.3. Effects of land-use on the initial litter mass loss

We used the data set from the temperate biome (228 sites out of 250; Table 1) to test the effect of land-use on litter mass loss. The model for land-use effects showed a significant interaction between land-use and tea type ($F = 41$, $P < .01$). However, post hoc contrasts showed no differences among land-use types for either Green or Rooibos tea (all comparisons: $P > .05$).

4. Discussion

The early stage of litter decomposition is a highly dynamic phase and therefore important for the understanding of litter decay and the controlling factors across biomes and ecosystem types. Here we studied the early stage mass loss of two standardized litter types (Green tea and Rooibos tea) across 336




Fig. 4. Relationship between remaining mass of Green tea and Rooibos tea and temperature (A) and precipitation (B) after the 3-month incubation period. Climatic variables were obtained from local weather stations or from wordclim for sites with no data. Circles show the mean values for each site and bars the standard errors. The regression line from the minimum adequate model is plotted only for the significant effects of precipitation and is obtained using only fixed factors. Band shows 95 confidence bands.




Fig. 5. A) Correlation between remaining mass of tea litter of different qualities (green and rooibos tea) after 3 month of incubation during the growing season. Symbols are arithmetic means for each biome and error bars indicate $\pm$ standard deviation. B) The average remaining mass aggregated by biome of Green tea (dashed line) and Rooibos tea (solid line) plotted against the mean annual precipitation for each biome (Table 1). The regression line is from a simple linear model showing significant effects for Green ($R^2 = 0.40$) and Rooibos ($R^2 = 0.64$).

sites globally and found that the litter type (quality) was the main determinant of the mass loss while climate and land use had little effect.

4.1. Substrate quality effects on litter decomposition

The effect of initial litter quality (chemical and physical composition) has been reported to be one of the key drivers of litter decomposition (Bradford et al., 2016; Cornwell et al., 2008; Heim and Frey, 2004). In our study, the litter type also had a strong control on initial decomposition as Green tea consistently decomposed faster than Rooibos tea (Fig. 3). Faster initial decomposition of Green tea is expected due to its higher fraction of water-soluble compounds in contrast to the low content of soluble or hydrolysable compounds in Rooibos tea (Didion et al., 2016). The mass loss of the litter during this early stage may be more related to the leaching losses than to microbial mineralization of soil organic C at the early stage of decomposition. In a pilot study, we measured changes in the initial weight after 3–4 min of cooking ($n = 332$) and recorded a weight loss of 31% for Green tea compared to 17% for Rooibos tea. Similar observation was made within different urban soil habitats by Pouyat et al. (2017). Moreover, Green and Rooibos tea differ in their carbon and nutrient chemistry (Keuskamp et al., 2013) and physical features (Didion et al., 2016). In a meta-analysis of the factors influencing mass loss rates involving 70 published studies, Zhang et al. (2008) demonstrated, similar to our study, the direct influence of litter

quality (C:N ratio and total nutrient content) on mass loss rates. The mass loss of both tea types decreased when precipitation increased (Table 2) which is in agreement with several studies showing a positive relationship between moisture availability and decomposition rates (Gholz et al., 2000; Prescott, 2010; García Palacios et al., 2016).

Overall, litter type explained 65% of the variability in litter mass loss at the global scale, which in turn implies that potential shifts in the relative abundance of vegetation types in the future caused by climatic changes could have large effects on global carbon budgets alone due to the differences in litter quality and consequently decomposition rates (Cornwell et al., 2008; Cornelissen et al., 2007).

4.2. Climate effects on litter mass loss

Across biomes, climatic factors are assumed to have a significant influence on litter decomposition by affecting the activity of decomposer organisms (Bradford et al., 2014); namely for every 10 °C increase in temperature a doubling of microbial decomposition is anticipated ($Q_{10} = 2$; Friedlingstein et al., 2006). Here, processes in the topsoil deserve special attention since they are particularly exposed to dynamic changes in environmental conditions.

We analyzed the across-site variation in initial litter mass loss at the site and biome scales. In this study, investigated sites are spread across large temperature and moisture gradients. We observed an effect of precipitation on early stage litter mass loss, while temperature did not show any significant effects (Fig. 3). Mean annual temperatures of <10 °C and moisture contents of <30% or >80% have been suggested as inhibiting thresholds for litter decay (Prescott, 2010). The absence of any significant effect of temperature on litter mass loss in our study may be a consequence of the fact that all sites incubated the tea bags during the “summer” under relatively favorable conditions where temperature values were generally within the “optimal” decay range. Furthermore, large variation in litter mass loss was observed for both litter types within any given biome (Fig. 5a, Table 2) suggesting that local-scale factors (e.g. soil properties, soil water content,

Table 3

Effects of climatic factors on the biome level remaining mass of the two tea types for data aggregated by biome (statistics relates to Fig. 5). Estimates obtained from simple linear models after backward selection. $R^2 = 0.84$.

	Est.(SE) ^a	t	P
Green tea	48.94(4.62)	10.60	<.01
Rooibos tea	88.23 (4.62)	19.10	<.01
PREC	−12.93(3.64)	−3.64	<.01

^a Models were fitted using precipitation/1000 to avoid very small estimates. Est. = estimates, SE = standard error.

disturbances) other than climate had strong controls on regional litter mass loss dynamics (Cornwell et al., 2008). Similarly, Ise and Moorcroft (2006) reported a low temperature sensitivity of decomposition ($Q_{10} = 1.37$) at the global scale. On the other hand, when examined separately, climate explained 40% of the variation for Green tea and 64% for Rooibos tea when the mean litter mass loss values were used for the given biome (Fig. 5b, Table 3). A similar finding was reported by Bradford et al. (2014), where the explanatory power of climate was increased to 84% when analyses were conducted on aggregated data.

Interestingly, early-stage litter mass losses of both litter types were comparable across all biomes (Fig. 3). The relative mass losses observed in the arctic sites may seem surprisingly high relative to the other warmer biomes. However, the study was carried out in the “summer season” where climatic conditions, even at the arctic sites are rather mild and warm and therefore favorable for decomposition (Couteaux et al., 1995). On the contrary, sites in the warmer biomes received less precipitation in the summer often being below potential evapotranspiration and leading to soil moisture deficit which again may result in lower mass losses. However, it has to be kept in mind that the results for arctic and arid-temperate biomes are based on a lower number of sites and should be interpreted with caution.

The data in this study collected during the growing season revealed that direct climatic control on early stage decomposition is of relatively minor importance. Instead, indirect climatic effects (e.g. plant community structure and associated microclimate, soil organic matter quality and structure of decomposer communities) may play a relatively stronger role in the early stage decomposition and may mask any importance of direct climatic controls (Aerts, 1997).

4.3. Land-use effects on litter mass loss

Long-term prevailing climatic conditions together with human activities define plant species composition and ecosystem structure, which in turn may affect decomposition rates. We did not observe any significant effects of land-use or management practices on the initial litter decomposition in the temperate biome. This may be caused by microbial decomposition not being limited by nutrients during the growing season. Another reason may be that in the early stage decomposition mineralization of labile C compounds is carried out by many groups of microorganisms while in the later stage of decomposition, decomposer groups may become more selected due to increased substrate complexity which in turn might lead to differences in litter mass loss between the land-use types (McGuire and Treseder, 2010). Hence, home-field advantage (Gholz et al., 2000) is expected to explain a fraction of the remaining variability at later and more advanced stages of decomposition. A detailed definition of different land-use categories would be necessary in order to be able to run more specific data analyses across all biomes.

5. Conclusions

Our study showed that litter type has the strongest influence on mass loss globally in the early stage of decomposition, while the effect of climate was only important under less favorable climatic conditions and when data were aggregated at the biome scale. This finding is particularly relevant for the general understanding of litter and carbon dynamics in relation to biosphere-atmosphere feedback, since the early stage litter decay is responsible for a significant fraction of the carbon

loss from litter, and because the lack of site specific climate control for this decomposition phase should be reflected in soil carbon models. The short-term period of just three month incubations used in this study provides insight into the short mass loss dynamics of plant litter. On the other hand the results cannot be extrapolated to capture a reliable signal of the long term nature of the decomposition rates, because long term decomposition involves other litter components and the drivers are likely to vary at spatial and temporal scales (Couteaux et al., 1995; Berg, 2014). Therefore caution should be payed when extrapolating from short-term to long-term rates (Moore et al., 2017). Therefore, the TeaComposition initiative includes additional sampling points after 12, 24, and 36 months, which will provide long term litter decomposition dynamics globally. Repeated observations over time (medium to long-term data) are essential for improving our understanding of the long term decay process of plant litter. Further, in addition to the observational networks included in this study (e.g. ILTER – see Mirtl et al., this issue), the TeaComposition initiative includes studies across collaborative experiments which are needed to identify and quantify the relative importance of multiple drivers (Verheyen et al., 2016; Borer et al., 2014).

Acknowledgements

This work was performed within the TeaComposition initiative, carried out by 190 institutions worldwide. We thank Gabrielle Drozdowski for her help with the packaging and shipping of tea, Zora Wessely and Johannes Spiegel for the creative implementation of the acknowledgement card, Josip Dusper for creative implementation of the graphical abstract, Christine Brendle for the GIS editing, and Marianne Debie for her help with the data cleaning. Further acknowledgements go to Adriana Principe, Melanie Köbel, Pedro Pinho, Margarida Santos-Reis, Thomas Parker, Steve Unger, Jon Gewirtzman and Margot McKleeven for the implementation of the study at their respective sites. We are very grateful to UNILEVER for sponsoring the Lipton tea bags and to the COST action ClimMani for scientific discussions, adoption and support to the idea of TeaComposition as a common metric. The initiative was supported by the following grants: ILTER Initiative Grant, ClimMani Short-Term Scientific Missions Grant (COST action ES1308; COST-STSM-ES1308-36004; COST-STSM-ES1308-39006; ES1308-231015-068365), INTERACT (EU H2020 Grant No. 730938), and Austrian Environment Agency (UBA). Franz Zehetner acknowledges the support granted by the Prometeo Project of Ecuador's Secretariat of Higher Education, Science, Technology and Innovation (SENESCYT). Ana I. Sousa, Ana I. Lillebø and Marta Lopes thanks for the financial support to CESAM (UID/AMB/50017/2013), to FCT/MEC through national funds, and the co-funding by the FEDER, within the PT2020 Partnership Agreement and Compete 2020. The research was also funded by the Portuguese Foundation for Science and Technology, FCT, through SFRH/BPD/107823/2015 (A.I. Sousa), co-funded by POPH/FSE.

Authorship

ID designed and coordinated the study with extensive input from CB, IKS, SKR, KSL accomplished data collection and preparation. SKR conducted statistical analyses. KV and BB provided inputs for manuscript concept. ID wrote the manuscript with contribution from all authors. The TeaComposition team implemented the study and provided site specific and climatic data. The authors declare no conflict of interest.

Appendix A

Table 2s

General characteristics of the study sites within the TeaComposition initiative.

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
4,01	Zöbelboden-IP1	Austria	47,836	14,443	950	6,9	1061	Temperate climate	Spruce forest, initial Cardamino trifoliae-Fagetum sensu Willner 2002	Ika Djukic
4,02	Zöbelboden-IP2	Austria	47,841	14,442	950	6,9	1061	Temperate climate	Mixed beech, spruce, maple, ash forest. Potential naural vegetation:Adenostylo glabrae-Fagetum sensu Willner 2002	Ika Djukic
4,03	Zöbelboden-IP3	Austria	47,839	14,444	950	6,9	1061	Temperate climate	Mixed spruce-beech forest	Ika Djukic
4,04	Zöbelboden-nutrient addition	Austria	47,841	14,435	950	6,9	1061	Temperate climate	Spruce forest; initial carbonate spruce-fir-beech forest	Ika Djukic
6	Klausen-Leopoldsdorf	Austria	48,108	16,075	510	8,1	724	Temperate climate	Beech forest	Ferdinand Kristöfel
7	Mondsee	Austria	47,881	13,350	860	7,2	1353	Temperate climate	Mixed spruce-broadleaved forest	Ferdinand Kristöfel
8	Mürzzuschlag	Austria	47,633	15,658	715	5,2	978	Temperate climate	Spruce forest	Ferdinand Kristöfel
9	Murau	Austria	47,061	14,111	1540	3,3	1366	Temperate climate	Spruce forest	Ferdinand Kristöfel
10	Jochberg	Austria	47,331	12,406	1050	3	1143	Temperate climate	Spruce forest	Ferdinand Kristöfel
11	AREC Raumberg-Gumpenstein	Austria	47,499	14,153	720	9,1	1088	Temperate climate	Meadow	Andreas Bohner
12	Neustift im Stubaital	Austria	47,117	11,300	970	6,5	852	Temperate climate	Managed grassland	Georg Wohlfahrt
13	Illmitz	Austria	47,767	16,750	113	10,1	599	Temperate climate	Managed grassland	Thomas Zechmeister
14	Pürgschachen Moor	Austria	47,581	14,346	632	7,3	1248	Temperate climate	Peat bog	Simon Drollinger
15	Jamtalferner	Austria	46,850	10,150	2960	-4,4	1374	Temperate climate	High alpine	Andrea Fischer
16	Jalhay-La Robinette	Belgium	50,550	6,067	500	7,7	1134	Temperate climate	Forest	Monique Carnol
17	Waroneu	Belgium	50,567	6,100	420	7,7	1134	Temperate climate	Forest	Monique Carnol
18	Brasschaat	Belgium	51,308	4,520	14	10	785	Temperate climate	Scots pine forest	Arne Verstraeten
19	Zoniënwood	Belgium	50,746	4,413	129	9,9	823	Temperate climate	Beech forest	Arne Verstraeten
20	Gontrode	Belgium	50,975	3,804	26	10	776	Temperate climate	Pedunculate oak - Beech forest	Arne Verstraeten
21	Ravels	Belgium	51,402	5,054	35	9,5	799	Temperate climate	Corsican pine forest	Arne Verstraeten
22	Wijnendale	Belgium	51,070	3,037	31	10,1	708	Temperate climate	Beech forest	Arne Verstraeten
23	Beklemeto	Bulgaria	42,783	24,609	1420	7,5	682	Temperate climate	Beech forest	Miglana Zhiyanski
24	Sofia-FRI	Bulgaria	42,630	23,353	650	8,6	602	Temperate climate	Cedrus atlantica trees	Maria Glushkova
25	Sofia-FRI	Bulgaria	42,631	23,204	650	8,6	581	Temperate climate	Grassland	Maria Glushkova
26	Govedarci	Bulgaria	42,233	23,439	1310	5,9	658	Temperate climate	Spruce forest	Miglana Zhiyanski
27	Govedarci	Bulgaria	42,238	23,438	1320	5,6	658	Temperate climate	Grassland	Miglana Zhiyanski
28	Mata dos Godoy State Park	Brazil	-23,433	-51,233	620	20,6	1486	Equatorial humid climate; tropical rain forest	Forest fragment and restoration site	Jose Marcelo Torezan
29	Congonhas Farm	Brazil	-22,733	-51,183	340	22,2	1285	Equatorial humid climate; tropical rain forest	Forest fragment and restoration site	Jose Marcelo Torezan
30	Alvorada Farm	Brazil	-22,983	-50,933	340	22	1271	Equatorial humid climate; tropical rain forest	Forest fragment and restoration site	Jose Marcelo Torezan
31,01	Natal Restinga Forest	Brazil	-5,896	-35,168	40	25,7	1298	Equatorial humid climate; tropical rain forest	Restinga forest	Adriano Caliman
31,02	Natal Restinga Shrubs	Brazil	-5,910	-35,179	50	25,7	1298	Equatorial humid climate; tropical rain forest	Restinga shrubland	Adriano Caliman
32,01	Bodoquena	Brazil	-20,986	-56,517	378	22,4	1353	Subtropical arid	Savannah forested	Franco Leandro de Souza
32,02	Bodoquena	Brazil	-20,998	-56,505	367	22,4	1353	Subtropical arid	Savannah forested	Franco Leandro de Souza
32,03	Bodoquena	Brazil	-21,000	-56,512	358	22,4	1353	Subtropical arid	Riparian forest	Franco Leandro de Souza
33,01	Restinga de Jurubatiba - Forest	Brazil	-22,264	-41,606	20	25,7	1298	Equatorial humid climate	Restinga forest	Rodrigo Lemes Martins
33,02	Restinga de Jurubatiba - Shrubs	Brazil	-22,264	-41,606	30	25,7	1298	Equatorial humid climate	Restinga shrubland	Rodrigo Lemes Martins
34,01	Floodplain Paraná River	Brazil	-22,799	-53,541	250	22,8	1280	Semi-arid tropical climate	Atlantic forest	Evanilde Benedito

(continued on next page)

Table 2s (continued)

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
34,02	Floodplain Paraná River	Brazil	-22,857	-53,600	250	22,8	1280	Semi-arid tropical climate	Atlantic forest and grassland	Evanilde Benedito
34,03	Floodplain Paraná River	Brazil	-22,721	-53,303	250	22,8	1280	Semi-arid tropical climate	Shrubland	Evanilde Benedito
34,04	Floodplain Paraná River	Brazil	-22,711	-53,276	250	22,8	1280	Semi-arid tropical climate	Shrubland and grassland	Evanilde Benedito
34,05	Floodplain Paraná River	Brazil	-22,774	-53,332	250	22,8	1280	Semi-arid tropical climate	Shrubland	Evanilde Benedito
34,06	Floodplain Paraná River	Brazil	-22,724	-53,218	250	22,8	1280	Semi-arid tropical climate	Atlantic forest	Evanilde Benedito
35	Tijuca National Park	Brazil	-22,963	-42,266	350	23	1157	Equatorial humid climate; tropical rain forest	NA	Vinicius Farjalla
36	Fazenda Miranda	Brazil	-15,731	-56,071	184	26	1268	Semi-arid tropical climate	Native forest	Francisco Lobo
37	Baia das Pedras	Brazil	-28,375	-68,276	127	26,2	1245	Subtropical arid climate	Native forest	Francisco Lobo
38	Parque Estadual do Utinga	Brazil	-1,431	-48,420	18	26,8	2369	Equatorial humid climate; tropical rain forest	NA	Thaísa Sala Michelan
39	Flashline Mars Arctic Research Station	Canada	75,431	-89,823	225	-17,3	131	Arctic climate	NA	Susan Holden Martin
40	Pan de Azúcar, fog zone	Chile	-26,150	-70,651	814	18	16	Subtropical arid climate	Desert with fog influence	Rafaella Canessa
41	Pan de Azúcar, interior zone	Chile	-26,150	-70,651	533	18	16	Subtropical arid climate	Desert	Rafaella Canessa
42	Reserva Quebrada de Talca	Chile	-30,011	-71,037	648	13,5	92	Mediterranean climate	Shrubland	Rafaella Canessa
43	Parque Nacional La Campana	Chile	-32,917	-71,150	726	13,4	377	Mediterranean climate	Mediterranean Forest	Rafaella Canessa
44	Parque Nacional Nahuelbuta	Chile	-37,783	-72,983	1205	8,1	1525	Temperate climate	Temperate Rain Forest	Rafaella Canessa
45	Monumento Nacional Contulmo	Chile	-38,017	-73,233	350	11	1544	Temperate climate	Temperate Rain Forest	Rafaella Canessa
46	Fray Jorge National Park	Chile	-30,667	-71,667	450	15,7	134	Temperate climate	Temperate Fog Forest	Aurora Gaxiola
47	LTSER Senda Darwin Biological Station	Chile	-42,467	-74,117	200	8,4	2140	Temperate climate	NA	Aurora Gaxiola
48	Punta Arenas	Chile	-53,167	-71,617	100	4,9	795	Temperate climate	Native Forest	Aurora Gaxiola
49	Fundo San Martín, Valdivia	Chile	-39,817	-73,150	115	11,7	2011	Temperate climate	Native Forest	Aurora Gaxiola
50	Omora Biosphere Reserve	Chile	-54,933	-67,317	50	4,7	480	Subantarctic climate	Native Forest	Aurora Gaxiola
51	Parque Nacional Nahuelbuta	Chile	-37,783	-72,983	1205	8,1	1525	Mediterranean climate	Temperate Rain Forest	Liesbeth van den Brink
52	Parque Nacional La Campana	Chile	-32,966	-71,084	721	13,4	377	Mediterranean climate	Mediterranean Forest	Liesbeth van den Brink
53	Reserva Quebrada de Talca	Chile	-30,011	-71,037	636	13,5	92	Mediterranean climate	Shrubland	Liesbeth van den Brink
54	Parque Nacional Pan de Azúcar	Chile	-26,150	-70,651	511	18	16	Subtropical arid climate	Desert	Liesbeth van den Brink
55	Hulunbeier grassland, Inner Mongolia	China	50,167	119,367	516	-1,8	374	Arid-temperate climate	Managed grassland	Wentao Luo
56,01	CATIE, Turrialba	Costa Rica	9,891	-83,648	600	22,4	3113	Equatorial humid climate; tropical rain forest	Mature secondary forest and mature disturbed forest	Geovana Carreno
56,02	CATIE, Turrialba	Costa Rica	9,896	-83,667	615	22,4	3113	Equatorial humid climate; tropical rain forest	Coffee agroforestry	Geovana Carreno
57	Nature Reserve Červený kříž	Czech Republic	49,993	13,931	420	7,5	584	Temperate climate	Oak forest	Petr Petřík
58	Bad Lauchstädt	Germany	51,395	11,876	119	9	492	Temperate climate	Grassland	Jutta Stadler
59	Bayreuth	Germany	49,973	11,514	336	7	720	Temperate climate	Deciduous Forest	Jutta Stadler
60,01	Rhine-Main-Observatory	Germany	50,154	9,002	115	9,5	665	Temperate climate	Grassland, intensively use	Marlen Mährlein
60,02	Rhine-Main-Observatory	Germany	50,174	9,064	115	9,5	662	Temperate climate	Grassland, intensively use	Marlen Mährlein
60,03	Rhine-Main-Observatory	Germany	50,132	8,965	130	9,9	644	Temperate climate	Deciduous Forest	Marlen Mährlein
60,04	Rhine-Main-Observatory	Germany	50,183	9,085	135	9,5	662	Temperate climate	Deciduous Forest	Marlen Mährlein
61	Landau	Germany	49,254	7,962	200	8,7	644	Temperate climate	Plot forest: mixed beech forest; vineyard; vineyard; stream floodplain: alluvial stream floodplain	Stefan Stoll
62	Hiddensee	Germany	54,550	13,100	1	8,2	545	Temperate climate	Coastal heath	Andrey Malyshev
63	Mols	Denmark	56,394	10,956	57	7,8	573	Temperate climate	Grass, heath	Inger Kappel Schmidt
64	Brandbjerg	Denmark	55,258	11,272	5	8,3	591	Temperate climate	Grass, heath	Klaus Steenberg Larsen
66	Odsherred	Denmark	55,833	11,700	30	8,2	602	Temperate climate	Forest	Inger Kappel Schmidt
67	Mattrup	Denmark	55,163	10,038	110	7,2	796	Temperate climate	Forest	Inger Kappel Schmidt
68	Valloe	Denmark	55,417	12,050	46	8,3	596	Temperate climate	NA	Inger Kappel Schmidt
69	Nørholm	Denmark	56,133	9,017	52	7,5	803	Temperate climate	NA	Inger Kappel Schmidt

70	Kragelund	Denmark	56,167	9,417	85	7,3	748	Temperate climate	NA	Inger Kappel Schmidt
71	Saarejärve-1	Estonia	58,657	26,757	56	4,9	606	Temperate climate	Pine forest	Ivika Ostonen
72	Saarejärve-2	Estonia	58,651	26,760	45	4,9	606	Temperate climate	Spruce forest	Ivika Ostonen
73	Vilsandi	Estonia	58,387	21,844	2	6	586	Temperate climate	Pine forest	Ivika Ostonen
74	Tõravere	Estonia	58,275	26,460	67	5	598	Temperate climate	Spruce forest	Ivika Ostonen
75	Sagadi	Estonia	59,562	26,046	45	4,8	624	Temperate climate	Pine forest	Ivika Ostonen
76	Vihula	Estonia	59,578	26,133	14	4,8	624	Temperate climate	Pine forest	Ivika Ostonen
77	Vändra	Estonia	58,708	25,064	43	5,2	672	Temperate climate	Spruce forest	Ivika Ostonen
78	Kuusnõmme	Estonia	58,307	21,971	5	6	592	Temperate climate	Mixed pine and spruce forest	Ivika Ostonen
79	Järvelja-1	Estonia	58,307	27,332	33	5	604	Temperate climate	Drained pine forest, monoculture	Ivika Ostonen
80	Järvelja-2	Estonia	58,303	27,287	31	5	604	Temperate climate	Drained spruce forest, monoculture	Ivika Ostonen
81	Järvelja-3	Estonia	58,289	27,316	33	5	604	Temperate climate	Drained birch forest	Ivika Ostonen
82	Lammi Biological Station	Finland	61,054	25,040	112	3,7	645	Boreal climate	Native broad-leaf and spruce forests	John Loehr
83	83c - Landemarais	France	49,002	-1,182	145	10,6	636	Temperate climate	Restored peatland	André-Jean Francez
84	Arboretum Champenoux, 54	France	48,751	6,338	256	9,4	765	Temperate climate	Exotic and local trees	Marie-Noëlle Pons
85	La Bouzule, 54	France	48,739	6,322	225	9,4	765	Temperate climate	Grassland	Marie-Noëlle Pons
86	Garden 1, Fléville-devant-Nancy	France	48,626	6,208	236	9,4	775	Temperate climate	Vegetable garden	Marie-Noëlle Pons
87	GISFI station, Homécourt, 54	France	49,222	6,003	231	9,5	795	Temperate climate	Afforested grassland	Florence Maunoury-Danger
88	Temperate Forest 1, Hémilly, 57	France	49,033	6,500	280	9,2	789	Temperate climate	Mixedforest	Florence Maunoury-Danger
89	Riparian forest, Liverdun, 54	France	48,756	6,058	200	9,3	743	Temperate climate	Alluvial forest	Michael Danger
90	Settling pond 1, Pompey, 54	France	48,769	6,136	207	9,3	743	Temperate climate	Afforested settling pond	Florence Maunoury-Danger
91	Settling pond 2, Russange, 54	France	49,483	5,931	378	8,9	818	Temperate climate	Afforested settling pond	Florence Maunoury-Danger
92	Gravel pit 1, Corny, 57	France	49,013	6,048	167	9,6	736	Temperate climate	Alluvial forest	Michael Danger
93	Gravel pit 2, Dieulouard, 54	France	48,829	6,084	177	9,3	743	Temperate climate	Alluvial forest	Michael Danger
94	Chitelet Botanical Garden, 88	France	48,042	7,003	1225	9,3	1344	Temperate climate	Wetland	Sylvie Dousset
95	JM Pelt Botanical Garden, 54	France	48,867	6,183	245	11,1	618	Temperate climate	Botanical garden	Sylvie Dousset
96	Forest soil SBL, Haye Forest, 54	France	48,639	6,122	382	11,1	618	Temperate climate	Mixed forest	Sylvie Dousset
97	Forest soil Rendzine, Haye Forest, 54	France	48,640	6,097	402	11,1	618	Temperate climate	Mixed forest	Sylvie Dousset
98	Haut Jacques - Podzol, 88	France	48,275	6,863	600	9,3	1344	Temperate climate	Mixed forest	Sylvie Dousset
99	Haut Jacques - SBA, 88	France	48,275	6,863	600	9,3	1344	Temperate climate	Mixed forest	Sylvie Dousset
100	Rudlin - SOP, 88	France	48,122	7,042	600	9,3	1344	Temperate climate	Alpine grassland	Sylvie Dousset
101	Rudlin - SBA, 88	France	48,122	7,042	600	9,3	1344	Temperate climate	Alpine grassland	Sylvie Dousset
108	LTZERZAA_ORCHAMP_CHAMROUSSE_1_CHAM1250	France	45,075	5,857	1249	7,5	1220	Temperate climate	Deciduous Broad-leaved Forest	Thomas Spiegelberger
109	LTZERZAA_ORCHAMP_CHAMROUSSE_2_CHAM1470	France	45,088	5,863	1471	7,5	1220	Temperate climate	Mixed Forest	Thomas Spiegelberger
110	LTZERZAA_ORCHAMP_CHAMROUSSE_3_CHAM1710	France	45,105	5,891	1713	6,2	1158	Temperate climate	Evergreen Coniferous Forest	Thomas Spiegelberger
111	LTZERZAA_ORCHAMP_CHAMROUSSE_4_CHAM1890	France	45,108	5,900	1887	4,7	1032	Temperate climate	Forest-grassland ecotone	Thomas Spiegelberger
112	LTZERZAA_ORCHAMP_CHAMROUSSE_5_CHAM2020	France	45,115	5,907	2021	4,7	1032	Temperate climate	Mountain Grassland	Thomas Spiegelberger
113	LTZERZAA_ORCHAMP_CHAMROUSSE_6_CHAM2180	France	45,122	5,914	2179	3,1	877	Temperate climate	Alpine meadow	Thomas Spiegelberger
118	LTZERZAA_ORCHAMP_RISTOLAS_1_RIS1870	France	44,746	6,999	1876	5,1	532	Temperate climate	Deciduous Coniferous Forest	Amélie Saillard
121	LTZERZAA_ORCHAMP_RISTOLAS_4_RIS2540	France	44,709	7,053	2555	1,75	403	Temperate climate	Alpine meadow	Amélie Saillard
125	FR AME CFE - Cime de Fer	France	44,326	6,938	2700	0,7	508	Temperate climate	Alpine meadow	Philippe Choler
129	FR AME LAU - Butte des Laussets	France	44,332	6,907	2508	2,5	674	Temperate climate	Subalpine grassland	Philippe Choler
132,01	Lyon (grasslands)	France	45,780	4,868	170	11,5	783	Temperate climate	Urban grassland	Pierre Marmonier
132,02	Lyon (undercover)	France	45,780	4,868	170	11,5	783	Temperate climate	Urban forest	Pierre Marmonier
133	Kerguelen Islands	France	-49,354	70,211	15	4,9	753	(Sub-)Arctic climate (Subantarctic climate)	Grassland	Marc Lebouvier
134	Forêt de Chaux	France	47,101	5,727	260	10,5	943	Temperate climate	Forest	Eric Lucot
135	Zone Atelier Plaine et Val de Sèvre	France	46,137	-0,490	66	12,4	901	Temperate climate	Agriculture	Vincent Bretagnolle
136	Tourbière de la Guette	France	47,322	2,281	165	11	705	Temperate climate	Peatland	Sébastien Gogo
137	Vosges (88)	France	48,168	5,942	420	9,2	852	Temperate climate	Agriculture	Marie-Noëlle Pons

(continued on next page)

Table 2s (continued)

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
138	Experimental station Gardouch	France	43,371	1,671	180	12,8	751	Temperate climate	Forest	Joël Merlet
139	Toulouse (VCG)	France	43,604	1,436	333	12,7	698	Temperate climate	Semi-natural grassland	Annie Ouin
140	Lüss	Germany	52,839	10,266	109	8,8	835	Temperate climate	Deciduous Forest	Meesenburg, Henning
141,01	Lange Bramke, Kamm	Germany	51,860	10,418	659	5,9	1339	Temperate climate	Coniferous forest	Meesenburg, Henning
141,02	Lange Bramke, Nordhang	Germany	51,855	10,413	597	5,9	1339	Temperate climate	Coniferous forest	Meesenburg, Henning
141,03	Lange Bramke, Südhang	Germany	51,856	10,413	597	5,9	1339	Temperate climate	Coniferous forest	Meesenburg, Henning
142	Solling, Buche	Germany	51,761	9,578	504	6,9	1193	Temperate climate	Deciduous Forest	Meesenburg, Henning
143	Solling, Fichte	Germany	51,765	9,580	508	6,9	1193	Temperate climate	Coniferous forest	Meesenburg, Henning
144	Göttinger Wald	Germany	51,528	10,047	420	8,4	773	Temperate climate	Deciduous Forest	Meesenburg, Henning
145	Augustendorf	Germany	52,908	7,859	33	9	820	Temperate climate	Deciduous Forest	Meesenburg, Henning
146	Ehrhorn	Germany	53,178	9,903	115	9	785	Temperate climate	Deciduous Forest	Meesenburg, Henning
147	Schafstaedt	Germany	51,374	11,736	172	8	611	Temperate climate	Meadow	Mark Frenzel
148	Friedeburg	Germany	51,604	11,721	98	8	611	Temperate climate	Meadow	Mark Frenzel
149	Greifenhagen	Germany	51,634	11,459	265	7,8	614	Temperate climate	Meadow	Mark Frenzel
150	Siptenfelde	Germany	51,655	11,049	397	7,8	561	Temperate climate	Pasture	Mark Frenzel
151	Harsleben	Germany	51,839	11,058	152	7,8	561	Temperate climate	Meadow	Mark Frenzel
152	Wanzleben	Germany	52,077	11,443	98	8,8	513	Temperate climate	Mown meadow	Mark Frenzel
153	Mueggelsee	Germany	52,000	14,683	35	9,1	553	Temperate climate	NA	Rita Adrian
154	Hohes Holz	Germany	52,087	11,222	193	8,8	529	Temperate climate	Mixed beech forest	Corinna Rebmann
155,01	Biodiversity-Exploratories, Hainich	Germany	50,967	10,400	330	7,3	702	Temperate climate	Mown pasture	Ute Hamer
155,02	Biodiversity-Exploratories, Hainich	Germany	51,083	10,417	330	7,3	702	Temperate climate	Mown pasture	Ute Hamer
155,03	Biodiversity-Exploratories, Hainich	Germany	51,100	10,417	330	7,7	666	Temperate climate	Mown pasture	Ute Hamer
155,04	Biodiversity-Exploratories, Hainich	Germany	51,267	10,417	330	7,7	666	Temperate climate	Mown pasture	Ute Hamer
155,05	Biodiversity-Exploratories, Hainich	Germany	51,267	10,317	330	7,7	695	Temperate climate	Mown pasture	Ute Hamer
155,06	Biodiversity-Exploratories, Hainich	Germany	51,050	10,383	330	7,3	778	Temperate climate	Mown pasture	Ute Hamer
155,07	Biodiversity-Exploratories, Hainich	Germany	51,000	10,400	330	7,7	666	Temperate climate	Mown pasture	Ute Hamer
155,08	Biodiversity-Exploratories, Hainich	Germany	51,200	10,417	330	7,3	702	Temperate climate	Pasture	Ute Hamer
155,09	Biodiversity-Exploratories, Hainich	Germany	51,017	10,367	330	7,7	695	Temperate climate	Pasture	Ute Hamer
155,1	Biodiversity-Exploratories, Hainich	Germany	51,269	10,433	330	7,7	666	Temperate climate	Mown pasture	Ute Hamer
155,11	Biodiversity-Exploratories, Hainich	Germany	51,267	10,450	330	7,7	695	Temperate climate	Mown pasture	Ute Hamer
155,12	Biodiversity-Exploratories, Hainich	Germany	51,300	10,367	330	7,7	695	Temperate climate	Mown pasture	Ute Hamer
155,13	Biodiversity-Exploratories, Hainich	Germany	50,967	10,750	330	7,7	695	Temperate climate	Pasture	Ute Hamer
155,14	Biodiversity-Exploratories, Hainich	Germany	51,283	10,367	330	7,8	580	Temperate climate	Pasture	Ute Hamer
155,15	Biodiversity-Exploratories, Hainich	Germany	51,217	10,583	330	7,7	695	Temperate climate	Meadow	Ute Hamer
155,16	Biodiversity-Exploratories, Hainich	Germany	51,267	10,500	330	7,9	638	Temperate climate	Meadow	Ute Hamer
155,17	Biodiversity-Exploratories, Hainich	Germany	51,267	10,500	330	7,9	638	Temperate climate	Meadow	Ute Hamer
155,18	Biodiversity-Exploratories, Hainich	Germany	51,067	10,433	330	7,9	638	Temperate climate	Mountain grassland	Ute Hamer
155,19	Biodiversity-Exploratories, Hainich	Germany	51,183	10,450	330	7,7	666	Temperate climate	Pasture	Ute Hamer
155,2	Biodiversity-Exploratories, Hainich	Germany	51,200	10,433	330	7,7	695	Temperate climate	Pasture	Ute Hamer
155,21	Biodiversity-Exploratories, Hainich	Germany	51,217	10,467	330	7,7	695	Temperate climate	Mown pasture	Ute Hamer
155,22	Biodiversity-Exploratories, Hainich	Germany	50,983	10,367	330	7,7	695	Temperate climate	Pasture	Ute Hamer
155,23	Biodiversity-Exploratories, Hainich	Germany	51,200	10,417	330	7,3	702	Temperate climate	Mown pasture	Ute Hamer
156,01	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,083	13,967	50	8,6	560	Temperate climate	Mown pasture	Ute Hamer
156,02	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,089	13,967	50	8,6	560	Temperate climate	Mown pasture	Ute Hamer
156,03	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	13,983	50	8,6	560	Temperate climate	Mown pasture	Ute Hamer
156,04	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,103	14,000	50	8,7	547	Temperate climate	Mown pasture	Ute Hamer
156,05	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	14,000	50	8,7	547	Temperate climate	Mown pasture	Ute Hamer
156,06	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	13,617	50	8,5	569	Temperate climate	Mown pasture	Ute Hamer
156,07	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,086	13,967	50	8,6	560	Temperate climate	Pasture	Ute Hamer
156,08	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	14,017	50	8,7	547	Temperate climate	Mown pasture	Ute Hamer
156,09	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,083	13,600	50	8,5	569	Temperate climate	Pasture	Ute Hamer

156,1	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,136	13,867	50	8,6	560	Temperate climate	Meadow	Ute Hamer
156,11	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	13,967	50	8,6	560	Temperate climate	Meadow	Ute Hamer
156,12	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	14,017	50	8,7	547	Temperate climate	Meadow	Ute Hamer
156,13	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,103	13,617	50	8,5	569	Temperate climate	Meadow	Ute Hamer
156,14	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,117	13,700	50	8,5	567	Temperate climate	Meadow	Ute Hamer
156,15	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,133	13,833	50	8,6	560	Temperate climate	Meadow	Ute Hamer
156,16	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,150	13,817	50	8,5	567	Temperate climate	Meadow	Ute Hamer
156,17	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	52,983	13,833	50	8,7	554	Temperate climate	Pasture	Ute Hamer
156,18	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	52,967	13,833	50	8,7	554	Temperate climate	Mown pasture	Ute Hamer
156,19	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,133	13,867	50	8,6	560	Temperate climate	Pasture	Ute Hamer
156,2	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	53,100	13,667	50	8,5	567	Temperate climate	Mown pasture	Ute Hamer
156,21	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	52,969	13,817	50	8,6	562	Temperate climate	Mown pasture	Ute Hamer
156,22	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	52,867	13,967	50	8,7	554	Temperate climate	Pasture	Ute Hamer
156,23	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	52,869	13,967	50	8,7	554	Temperate climate	Pasture	Ute Hamer
156,24	Biodiversity-Exploratories, Schorfheide-Chorin	Germany	52,967	13,817	50	8,6	562	Temperate climate	Pasture	Ute Hamer
157,01	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,333	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,02	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,367	9,467	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,03	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,400	9,517	730	7,1	923	Temperate climate	Meadow	Ute Hamer
157,04	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,367	9,417	730	7,5	911	Temperate climate	Mown pasture	Ute Hamer
157,05	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,436	730	7,5	911	Temperate climate	Mown pasture	Ute Hamer
157,06	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,400	9,433	730	7,5	911	Temperate climate	Mown pasture	Ute Hamer
157,07	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,367	730	7,5	911	Temperate climate	Pasture	Ute Hamer
157,08	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,417	9,483	730	7,5	911	Temperate climate	Pasture	Ute Hamer
157,09	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,500	730	7,1	923	Temperate climate	Pasture	Ute Hamer
157,1	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,367	9,200	730	7,8	905	Temperate climate	Meadow	Ute Hamer
157,11	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,483	9,433	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,12	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,483	9,433	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,13	Biodiversity-Exploratories, Schwäbische Alb	Germany	44,367	9,517	730	9,7	942	Temperate climate	Mown pasture	Ute Hamer
157,14	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,400	9,500	730	7,1	923	Temperate climate	Mown pasture	Ute Hamer
157,15	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,400	730	7,5	911	Temperate climate	Pasture	Ute Hamer
157,16	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,450	9,450	730	7,5	911	Temperate climate	Mown pasture	Ute Hamer
157,17	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,450	9,453	730	7,5	911	Temperate climate	Mown pasture	Ute Hamer
157,18	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,450	9,483	730	7,5	911	Temperate climate	Pasture	Ute Hamer
157,19	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,433	9,417	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,2	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,417	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,21	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,400	9,450	730	7,5	911	Temperate climate	Meadow	Ute Hamer
157,22	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,383	9,433	730	7,5	911	Temperate climate	Pasture	Ute Hamer
157,23	Biodiversity-Exploratories, Schwäbische Alb	Germany	48,450	9,500	730	7,1	923	Temperate climate	Pasture	Ute Hamer
158	Síkfőkút Project	Hungary	47,917	20,433	345	9,4	565	Temperate climate	Deciduous Forest	Zsolt Kotroczó and István Fekete
159	Kiskunság LTER - Fülöpháza	Hungary	47,453	19,704	108	10,6	522	Temperate climate	Grassland	Erzsébet Hornung
160	Shita	Israel	30,147	35,120	250	19,4	207	Subtropical arid climate	Desert	Elli Groner
161	Ramon	Israel	31,250	35,369	440	21,3	60	Subtropical arid climate	Desert	Elli Groner
162,01	Matsch-Mazia	Italy	46,677	10,576	1000	1,6	528	Temperate climate	Dry pasture	Julia Seeber
162,02	Matsch-Mazia	Italy	46,684	10,585	1500	1,6	528	Temperate climate	Dry pasture	Julia Seeber
162,03	Matsch-Mazia	Italy	46,692	10,593	2000	1,6	528	Temperate climate	Dry pasture	Julia Seeber
162,04	Matsch-Mazia	Italy	46,703	10,595	2500	1,6	528	Temperate climate	Dry pasture	Julia Seeber
163	Kanumazawa Riparian Research Forest	Japan	39,100	141,850	450	9,2	2056	Temperate climate	Forest, deciduous	Kazuhiko Hoshizaki
164,01	University of Tokyo Chichibu Forest	Japan	35,919	138,831	880	9	1554	Temperate climate	Natural forest	Satoshi Suzuki
164,02	University of Tokyo Chichibu Forest	Japan	35,917	138,818	1320	6,6	1554	Temperate climate	Natural forest	Satoshi Suzuki
164,03	University of Tokyo Chichibu Forest	Japan	35,915	138,801	1780	3,6	1554	Temperate climate	Natural forest	Satoshi Suzuki
165	Kasuya Research Forest	Japan	33,653	130,545	520	14,6	1917	Warm-temperate, humid climate	Natural forest	Tsutomu Enoki
166	Uryu	Japan	44,356	142,258	300	4,4	1400	Temperate climate	Cool temperate mixed forest (evergreen coniferous and deciduous broad-leaved species)	Hideaki Shibata
167	Yamashiro Experimental Forest	Japan	34,783	135,850	255	13,8	1676	Warm-temperate, humid climate	Secondary forest, deciduous	Karibu Fukuzawa
168	Ashoro	Japan	43,263	143,508	330	5,5	1051	Temperate climate	Forest, deciduous	Mioko Ataka, Yuji Kominami
169	Akazu Research Forest	Japan	36,217	137,167	304	9,7	1838	Temperate climate	Secondary forest, deciduous	Yasuhiro Utsumi
170,01	Mt. Hakkoda Forest_400B	Japan	40,593	140,964	416	9	1501	Warm-temperate, humid climate	Natural forest	Takanori Sato
170,02	Mt. Hakkoda Forest_600B	Japan	40,596	140,946	649	7,9	1501	Warm-temperate, humid climate	Natural forest	Hiroko Kurokawa
170,03	Mt. Hakkoda Forest_800B	Japan	40,636	140,931	791	7	1501	Warm-temperate, humid climate	Natural forest	Hiroko Kurokawa
170,04	Mt. Hakkoda Forest_1000A	Japan	40,660	140,851	980	6,5	1501	Warm-temperate, humid climate	Natural forest	Hiroko Kurokawa

(continued on next page)

Table 2s (continued)

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
170,05	Mt. Hakkoda Forest_1200A	Japan	40,666	140,867	1214	5,5	1501	Warm-temperate, humid climate	Natural forest	Hiroko Kurokawa
170,06	Mt. Hakkoda Forest_1400A	Japan	40,673	140,874	1404	4,9	1412	Warm-temperate, humid climate	Natural forest	Hiroko Kurokawa
171	Ashiu Experimental Forest	Japan	36,010	137,003	260	9	2065	Temperate climate	Natural Forest	Takeshi Ise
172	Kamigamo Experimental Station	Japan	34,083	135,767	220	12,3	2498	Warm-temperate, humid climate	Natural Forest	Naoko Tokuchi
173	Shiiba Research Forest	Japan	32,398	131,173	1050	12,5	3072	Warm-temperate, humid climate	Natural mixed forest	Takuo Hishi
174	Sugadaira	Japan	36,523	138,500	1320	10,6	1239	Warm-temperate, humid climate	Grassland, Natural forest	Tanaka Kenta
175	Tomakomai Experimental Forest	Japan	42,699	141,571	80	6,7	1112	Temperate climate	Secondary forest, deciduous	Tatsuro Nakaji
176	Engure LTSER	Latvia	57,289	23,154	10	6,3	634	Temperate climate	Pine Forest	Tsutom Hiura
179	Engure LTSER	Latvia	57,302	23,049	7	6,3	634	Temperate climate	Deciduous Forest	Inara Melece
180,01	Aukstaitija IMS	Lithuania	55,464	26,004	188	5,7	658	Temperate climate	Forest, coniferous	Inara Melece
180,02	Aukstaitija IMS	Lithuania	55,453	26,068	159	5,7	658	Temperate climate	Forest, coniferous	Algirdas
181	Zemaitija IMS	Lithuania	56,017	21,887	170	6,1	790	Temperate climate	Forest, coniferous	Augustaitis
182	Forest Research Institute Malaysia, Kepong	Malaysia, Selangor	3,235	101,633	82	26,1	358	Equatorial humid climate; tropical rain forest	Planted and naturally regenerating forest	Algirdas
183	SSDE-1	Mali	15,317	-9,050	270	28,1	1500	Subtropical arid climate	NA	Jeyanny
184	SSDE-2	Mali	14,533	-9,967	262	27,9	712	Semi-arid tropical climate	NA	Vijayanathan
185	SSDE-3	Mali	12,883	-8,483	370	27	986	Semi-arid tropical climate	NA	Niall Hanan
186	SSDE-4	Mali	11,600	-7,050	368	27,2	1017	Semi-arid tropical climate	NA	Niall Hanan
187	SSDE-5	Mali	11,033	-6,083	347	27,1	1105	Semi-arid tropical climate	NA	Niall Hanan
188	Estero Pargo	Mexico	18,651	-91,759	1	26,4	1502	Equatorial humid climate; tropical rain forest	Natural mangrove forest	José Gilberto Cardoso
189	ESTERO DE URIAS LAGOON	Mexico	23,173	-106,326	1	24,8	752	Mediterranean climate	Natural mangrove forest	Mohedano
192	MARISMAS NACIONALES	Mexico	22,410	-105,636	1	25,1	1627	Mediterranean climate	Natural mangrove forest	Ana Carolina Ruiz Fernández
193	SALAZAR FOREST	Mexico	19,286	-99,383	3124	12,4	1098	Subtropical arid climate	Sacred fir and pinus forest	Joan Albert
201	Wadi Nar station	Palestine	31,724	35,286	415	18,3	412	Subtropical arid climate	Olive orchard	Sánchez Cabeza
202,01	Companhia das Lezírias	Portugal	38,843	-8,765	60	17,4	774	Mediterranean climate	Evergreen cork oak forest	Eduardo Ordoñez Regil
202,02	Companhia das Lezírias	Portugal	38,851	-8,782	43	17,4	774	Mediterranean climate	Evergreen cork oak forest	Jawad Shoqeir
202,03	Companhia das Lezírias	Portugal	38,857	-8,783	47	17,4	774	Mediterranean climate	Evergreen cork oak forest	Cristina
202,04	Companhia das Lezírias	Portugal	38,834	-8,809	43	17,4	774	Mediterranean climate	Evergreen cork oak forest	Branquinho
202,05	Companhia das Lezírias	Portugal	38,826	-8,813	42	17,4	774	Mediterranean climate	Evergreen cork oak forest	Cristina
202,06	Companhia das Lezírias	Portugal	38,814	-8,801	50	17,4	774	Mediterranean climate	Evergreen cork oak forest	Branquinho
202,07	Companhia das Lezírias	Portugal	38,804	-8,816	45	17,4	774	Mediterranean climate	Evergreen cork oak forest	Cristina
202,08	Companhia das Lezírias	Portugal	38,835	-8,818	28	17,4	774	Mediterranean climate	Evergreen cork oak forest	Branquinho
202,09	Companhia das Lezírias	Portugal	38,837	-8,835	27	17,4	774	Mediterranean climate	Evergreen cork oak forest	Cristina
202,1	Companhia das Lezírias	Portugal	38,827	-8,840	30	17,4	774	Mediterranean climate	Evergreen cork oak forest	Branquinho
202,11	Companhia das Lezírias	Portugal	38,811	-8,849	31	17,4	774	Mediterranean climate	Evergreen cork oak forest	Cristina
202,12	Companhia das Lezírias	Portugal	38,818	-8,855	28	17,4	774	Mediterranean climate	Evergreen cork oak forest	Branquinho
203	Ria de Aveiro	Portugal	40,602	-8,740	1	14,3	800	Mediterranean climate	Wetland, Salt marsh	Cristina
203,01	Ria de Aveiro	Portugal	40,601	-8,741	1	14,3	800	Mediterranean climate	Wetland, Salt marsh	Branquinho
204	Luquillo Experimental Forest	Puerto Rico	18,344	-65,826	61	25,4	1943	Equatorial humid climate;	NA	Ana I. Lillebø
										Jill Thompson

205	Elevational gradient	Puerto Rico	18,344	-65,826	61	25,1	2003	tropical rain forest Equatorial humid climate; tropical rain forest	NA	Grizelle González
206	Braila Islands LTSE	Romania	44,885	27,861	9	11,5	454	Arid-temperate climate	Wetland	Elena Preda
207	Braila Islands LTSE	Romania	44,885	27,861	9	11,5	454	Arid-temperate climate	Wetland	Elena Preda
208	Neajlov basin LTSE	Romania	44,340	25,667	85	10,8	598	Temperate climate	Forest	Elena Preda
209	Neajlov basin LTSE	Romania	44,340	25,667	85	10,8	598	Temperate climate	Forest	Elena Preda
210,01	Fruska gora	Serbia	45,135	19,642	403	11,1	679	Temperate climate	Deciduous Forest	Dušanica Krašić
210,02	Fruska gora	Serbia	45,138	19,647	478	11,1	679	Temperate climate	Deciduous Forest	Dušanica Krašić
210,03	Fruska gora	Serbia	45,137	19,676	468	11,1	679	Temperate climate	Deciduous Forest	Dušanica Krašić
212,01	Podunajská nížina Lowland forest	Slovakia	48,277	17,321	173	9,4	669	Temperate climate	Vineyard on loess	Róbert Kanka
212,02	Podunajská nížina Lowland vineyard	Slovakia	48,276	17,321	173	9,4	669	Temperate climate	Pannonian oak and hornbeam forest	Róbert Kanka
212,03	Podunajská nížina Lowland grove grassland	Slovakia	48,306	17,287	177	9,4	669	Temperate climate	Cherry orchard (Cerasus avium)	Róbert Kanka
212,04	Podunajská nížina Lowland orchard-garden	Slovakia	48,306	17,287	177	9,4	669	Temperate climate	Lowland ruderalised meadow	Róbert Kanka
213	Tatry, LTER	Slovakia	49,083	20,233	1100	5,4	781	Temperate climate	Temperate oniferous forest	Peter Fleischer
214	Kralova hola	Slovakia	48,887	20,128	1850	3,8	1017	Temperate climate	Alpine grassland	Veronika Piscová
215	Jalovecka dolina	Slovakia	49,218	19,672	1893	2,9	1259	Temperate climate	Alpine grassland	Veronika Piscová
216	Báb	Slovakia	48,303	17,889	190	9,7	600	Temperate climate	Thermophilic oak forest	Veronika Piscová
217	Kremnicke vrchy Ecological Experimental Station	Slovakia	48,635	19,070	500	7,8	742	Temperate climate	Temperate deciduous forest	Milan Barna
218	Hodruska vrchovina	Slovakia	48,550	18,858	470	7,6	768	Temperate climate	Temperate deciduous forest	Milan Barna
219	Stiavnicke vrchy	Slovakia	48,552	18,947	600	7,6	768	Temperate climate	Temperate deciduous forest	Milan Barna
220	Javorie	Slovakia	48,504	19,188	785	6,7	794	Temperate climate	Temperate deciduous forest	Milan Barna
222	Wolwekraal Nature Reserve	South Africa	-33,197	22,029	567	7,8	177	Subtropical arid climate	Protected Nature Reserve	Joh Henschel
223	Tierberg Karoo Research Station, SAEON Arid Lands Node	South Africa	-33,165	22,268	752	17,8	177	Subtropical arid climate	Livestock/large game enclosure within wildlife ranch	Joh Henschel
224	Collserola	Spain	41,430	2,082	255	16,1	613	Mediterranean climate	Protected Nature Reserve	Anna Avila
225	Montseny	Spain	41,467	2,210	760	12,6	839	Mediterranean climate	Protected Nature Reserve	Fernando Maestre
226	Valdemoro	Spain	40,190	-3,601	622	16,6	631	Mediterranean climate	Protected area with wild and domestic grazers	Fernando Maestre
228	Aneboda IM	Sweden	57,114	14,551	240	5,8	750	Temperate climate	Coniferous forest	Stefan Löfgren
229	Kindla IM	Sweden	59,754	14,908	320	4,2	900	Boreal climate	Coniferous forest	Stefan Löfgren
230	Vordemwald	Switzerland	47,274	7,887	480	8,8	1028	Temperate climate	Temperate mixed forest	Marcus Schaub
231	Bettlachstock	Switzerland	47,225	7,417	1149	7,4	1113	Temperate climate	Temperate deciduous forest	Marcus Schaub
232	Pfynwald	Switzerland	46,303	7,612	615	3,6	1418	Temperate climate	Xeric mature Scots pine forest	Marcus Schaub
233	Novaggio	Switzerland	46,023	8,840	950	9,9	1272	Temperate climate	Unmanaged former coppice forest	Marcus Schaub
234	Beatenberg	Switzerland	46,700	7,762	1511	6,2	1235	Temperate climate	Temperate spruce forest	Marcus Schaub
235	Schänis	Switzerland	47,165	9,067	733	6	1364	Temperate climate	Temperate beech forest	Marcus Schaub
236	Birmensdorf	Switzerland	47,362	8,454	550	8,8	1103	Temperate climate	Temperate mixed forest	Marcus Schaub
237	Salgesch	Switzerland	46,317	7,583	805	3,6	1418	Temperate climate	Xeric mature Scots pine forest	Marcus Schaub
238	Fushan	Taiwan	24,759	121,598	720	21	3025	Warm-temperate, humid climate	Natural subtrpical mixed broadleaf rain forest	Chiao-Ping Wang
239	YYL	Taiwan	24,590	121,416	1650	15,1	2659	Warm-temperate, humid climate	Subtropical mountain cloud coniferous forest	Chiao-Ping Wang
240	12 experimental sites	UK	0,000	0,000	NA	NA	NA	Temperate climate	NA	Jill Thompson
241	Harvard Forest	USA	42,000	-73,200	310	7,3	1246	Temperate climate	Temperate forest	Jim Tang
242	Toolik Station	USA	68,633	-149,600	760	-11,7	229	Arctic climate	Arctic tundra	Jim Tang
243	Waquoit Bay salt marsh	USA	41,367	-70,500	1	10	1138	Temperate climate	Salt marsh	Jim Tang
244	H.J. Andrews Forest	USA	44,367	122,367	162	7,9	1663	Temperate climate	Old-growth forest	Kate Lajtha
245	Central Arizona-Phoenix	USA	33,604	-112,498	448	21,1	198	Arid-temperate climate	Desert	Sally Wittlinger
246	Mansfield_SC1	USA	44,507	-72,836	565	5,2	1070	Temperate climate	Mixed forest	Carol Adair
247	Smithsonian Environmental Research Center	USA	38,883	-76,550	100	13,3	1091	Temperate climate	Deciduous forest	Katalin Szlavecz
248	Smithsonian Global Change Research Wetland	USA	38,889	-77,026	2	12,9	1035	Temperate climate	Salt marsh	Thomas J. Mozdzer
249	PIE-LTER (TIDE Project)	USA	42,721	70,848	1	9,5	1191	Temperate climate	Salt marsh	Thomas J. Mozdzer
250	Reynolds Creek CZO	USA	43,205	-116,750	1200	7,7	330	Arid-temperate climate	Sagebrush steppe	Marie-Anne de Graaff
251	Cedar Point Biological Station	USA	41,207	-101,667	982	9,1	447	Arid-temperate climate	Short Grass Prairie	Johannes M H Knops
252,01	Bartlett Experimental Forest Site C6	USA	44,040	-71,275	460	5,5	1270	Temperate climate	Northern hardwood forest	Ruth Yanai
252,02	Bartlett Experimental Forest Site C8	USA	44,054	-71,298	330	5,5	1270	Temperate climate	Northern hardwood forest	Ruth Yanai
253	Hubbard Brook Experimental Forest (MELNHE)	USA	43,933	-71,733	500	7,4	1123	Temperate climate	Northern hardwood forest	Matt Vadeboncoeur
254	Jeffers Brook	USA	44,050	-72,467	730	5,1	1077	Temperate climate	Northern hardwood forest	Ruth Yanai
255	Hubbard Brook Experimental Forest (ISE)	USA	43,936	-71,758	500	7,4	1123	Temperate climate	Northern hardwood forest	Matt Vadeboncoeur

(continued on next page)

Table 2s (continued)

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
256	Hubbard Brook Experimental Forest (DroughtNet)	USA	43,946	-71,701	265	7,4	1123	Temperate climate	Northern hardwood forest	Matt Vadeboncoeur
258	Cummins Creek Wilderness Area, Oregon	USA	44,450	-124,167	NA	9,4	2555	Temperate climate	NA	Andy Moldenke
259	Mary's Peak, Oregon	USA	44,833	-123,933	98	10,4	2215	Temperate climate	NA	Andy Moldenke
260	Andrews Forest, LTER, Oregon	USA	44,367	-122,417	564	8,6	2072	Temperate climate	NA	Andy Moldenke
261	Andrews Forest, LTER, Oregon	USA	44,367	-122,217	628	6,8	2143	Temperate climate	NA	Andy Moldenke
262	Andrews Forest, LTER, Oregon	USA	44,367	-122,217	628,000	6,800	2143,000	Temperate climate	NA	Andy Moldenke
263	Metolius River Natural Area, Oregon	USA	44,817	-122,050	739,000	7,100	2123,000	Temperate climate	NA	Andy Moldenke
264	Sisters, Oregon	USA	44,291	-121,549	971,000	6,600	641,000	Temperate climate	NA	Andy Moldenke
265	Sky Oaks Field Station	USA	33,350	116,633	1420	15,4	269	Mediterranean climate	Chaparral	George Vourlitis
266	Santa Margarita Ecological Reserve	USA	33,483	117,181	254	16,6	396	Mediterranean climate	Coastal sage scrub (soft chaparral)	George Vourlitis
267	Ten Thousand Islands National Wildlife Refuge	USA	25,233	-81,117	0	23,8	1219	Semi-arid tropical climate	NA	Sean Charles
272,01	Tigirek Strict Reserve, Plot 01	Russia	51,057	82,987	1426	1,6	1120	Temperate climate	Alpine meadow	Evgeny Davydov
272,03	Tigirek Strict Reserve, Plot 03	Russia	51,110	83,052	994	1,6	980	Temperate climate	Meadow	Evgeny Davydov
272,05	Tigirek Strict Reserve, Plot 05	Russia	51,051	82,975	1493	1,6	1120	Temperate climate	Natural forest (Pinus sibirica open forest)	Evgeny Davydov
272,06	Tigirek Strict Reserve, Plot 06	Russia	51,045	83,000	1572	1,6	1120	Temperate climate	Natural forest + meadow (timberline)	Evgeny Davydov
272,07	Tigirek Strict Reserve, Plot 07	Russia	51,040	82,998	1391	1,6	1120	Temperate climate	Natural forest (montane)	Evgeny Davydov
272,08	Tigirek Strict Reserve, Plot 08	Russia	51,041	82,999	1453	1,6	1120	Temperate climate	Natural forest + meadow (timberline)	Evgeny Davydov
272,09	Tigirek Strict Reserve, Plot 09	Russia	51,010	82,999	1537	1,6	1120	Temperate climate	Natural forest + meadow (timberline)	Evgeny Davydov
272,1	Tigirek Strict Reserve, Plot 10	Russia	51,115	83,016	948	1,6	980	Temperate climate	Natural forest (Abies sibirica)	Evgeny Davydov
272,12	Tigirek Strict Reserve, Plot 12	Russia	51,045	82,989	1526	1,6	1120	Temperate climate	Natural forest (Pinus sibirica open forest)	Evgeny Davydov
272,13	Tigirek Strict Reserve, Plot 13	Russia	51,056	82,987	1455	1,6	1120	Temperate climate	Subalpine tall-grasses	Evgeny Davydov
272,14	Tigirek Strict Reserve, Plot 14	Russia	51,057	82,987	1432	1,6	1120	Temperate climate	Alpine meadow	Evgeny Davydov
273,01	State Nature Reserv "Stolby", Plot 01	Russia	55,912	92,733	703	1,1	552	Boreal climate	Natural forest (Pinus sylvestris L., Larix sibirica Ledeb.)	Elena Tropina
273,02	State Nature Reserv "Stolby", Plot 02	Russia	55,946	92,825	285	1,2	471	Boreal climate	Natural forest (Populus tremula L.)	Elena Tropina
273,03	State Nature Reserv "Stolby", Plot 03	Russia	55,707	92,934	239	1,2	471	Boreal climate	Natural forest (Betula pendula Roth)	Elena Tropina
273,04	State Nature Reserv "Stolby", Plot 04	Russia	55,737	92,782	218	1,2	471	Boreal climate	Mesophytic meadow	Elena Tropina
273,05	State Nature Reserv "Stolby", Plot 05	Russia	55,785	92,722	214	1,2	471	Boreal climate	Mesophytic meadow	Elena Tropina
273,06	State Nature Reserv "Stolby", Plot 06	Russia	55,827	92,811	722	1,1	552	Boreal climate	Natural forest (Pinus sylvestris L., Larix sibirica Ledeb.)	Elena Tropina
273,07	State Nature Reserv "Stolby", Plot 07	Russia	55,845	92,833	673	1,1	552	Boreal climate	Natural forest (Abies sibirica Ledeb.) + wet meadow	Elena Tropina
273,08	State Nature Reserv "Stolby", Plot 08	Russia	55,912	92,886	208	1,2	471	Boreal climate	Mesophytic meadow	Elena Tropina
273,09	State Nature Reserv "Stolby", Plot 09	Russia	55,867	92,936	709	1,1	552	Boreal climate	Natural forest (Pinus sylvestris L., Larix sibirica Ledeb., Populus tremula L.)	Elena Tropina
273,10	State Nature Reserv "Stolby", Plot 10	Russia	55,891	92,924	263	1,2	471	Boreal climate	Mesophytic meadow	Elena Tropina
274	State Nature Reserv "Olekminsky"	Russia	58,000	121,000	450	-8,6	424	Boreal climate	Natural forest(Pinus sylvestris L., Larix gmelinii(Rupr.) Rupr.)	Yury Rozhkov
275	Gossenköllesee	Austria	47,229	11,014	2417	3,2	1112	Temperate climate	High alpine	Birgit Sattler
278	Eight Mile Lake, Healy, Alaska	USA	63,876	-149,247	684	-1	384	Boreal climate	Boreal-tundra ecotone	Rebecca Hewitt
279	Murphy Dome, Fairbanks, Alaska	USA	64,882	-148,391	210	-3	275	Boreal climate	Boreal forest	Rebecca Hewitt
280	VCU_Rice_Rivers_Center_Swamp	USA	37,327	-77,208	0	14,3	1123	Temperate climate	Tidal Swamp Wetland	Joe Morina
281	Xishuangbanna	China	22,010	100,800	556	21,7	1460	Semi-arid tropical climate	Primary forest	Wenjun Zhou
282	Yuanjiang	China	28,945	112,598	30	24,3	790	Warm-temperate, humid climate	Savannah forested	Wenjun Zhou
283	Ailao Mountain	China	23,833	101,567	1852	11	1980	Semi-arid tropical climate	Primary forest	Wenjun Zhou
284	Lijiang	China	26,865	100,229	2517	9,1	1160	Arid-temperate climate	Primary forest	Wenjun Zhou
285	Jilin	China	42,383	128,083	802	2,5	688	Temperate climate	Secondary forest and white birch plantation	Yalin Hu
286	Liaoning	China	41,847	124,934	597	4,8	885	Temperate climate	Laruch monoculture	Yalin Hu
287	Zhejiang	China	29,967	122,350	786	16,7	1249	Warm-temperate, humid climate	Secondary forest and chinese fir plantation	Yalin Hu
288	Fujian	China	26,557	118,112	360	18,7	1729	Semi-arid tropical climate (Subtropical climate)	Secondary forest and chinese fir plantation	Yalin Hu
289	Hainan	China	18,730	108,890	800	21,7	1523	Semi-arid tropical climate (Tropical climate)	Secondary forest and chinese fir plantation	Yalin Hu
290	Jiangxi	China	24,562	114,431	550-600	18,5	1821	Semi-arid tropical climate (Subtropical climate)	Secondary forest and chinese fir plantation	Yalin Hu
291	Hunan	China	26,849	109,606	432	16,5	1280	Semi-arid tropical climate (Subtropical climate)	Secondary forest and chinese fir plantation	Yalin Hu
292	Inner Mongolia	China	42,500	122,317	120	7,6	506	Arid-temperate climate	Mongolian pine monoculture	Wentao Luo
293	Cattai, NSW, Lilly	Australia	-31,829	152,639	5	14,5	799	Warm-temperate, humid climate	Restored swamp	Stacey Trevathan-Tackett
294	Cattai, NSW, Melaluca	Australia	-31,830	152,637	11	14,5	799	Warm-temperate, humid climate	Restored swamp	Stacey Trevathan-Tackett

295	Darawakh, NSW	Australia	-32,091	152,488	3	14,5	799	Warm-temperate, humid climate	Seasonal wetland	Stacey Trevathan-Tackett
296	Rhyll, Victoria	Australia	-38,457	145,290	0	14,3	832	Temperate climate	Grassland	Stacey Trevathan-Tackett
297	Rhyll, Victoria	Australia	-38,457	145,289	0	14,3	832	Temperate climate	Mangrove	Stacey Trevathan-Tackett
298	Rhyll, Victoria	Australia	-38,459	145,288	0	14,3	832	Temperate climate	Succulent saltmarsh	Stacey Trevathan-Tackett
301	Northeast Science Station, Cherskiy, Russia	Russia	68,743	161,407	30	-11,6	230	Arctic climate	Larch forest	Heather Alexander
302	Mukhrino Field Station	Russia	60,889	68,703	50	8,2	545	Boreal climate	Raised bog	Nina Filippova
303	Hiddensee	Germany	54,551	13,104	1	-0,9	536	Temperate climate	Heathland	Andrey Malyshev
305	Hanshagen	Germany	54,054	13,514	45	8,3	562	Temperate climate	Beech forest	Robert Weigel
317	Aktru	Russia	50,083	87,782	2140	-5,2	430	Boreal climate	Alpine tundra	Roberto Cazzolla Gatti
318	Ob River	Russia	57,200	84,317	70	0,3	532	Boreal climate	Taiga forest and wetlands	Roberto Cazzolla Gatti
319	Mediterranean Shrublands	Italy	40,757	16,913	348	13,6	650	Mediterranean climate	Oak forests and shrubland	Roberto Cazzolla Gatti
320	Igloolik (Nunavut)	Canada	69,398	-81,543	15	-14,4	115	Arctic climate	Tundra	Nicolas Lecomte
321	Fendt	Germany	48,375	11,108	600	8,7	982	Temperate climate	Grassland	Ralf Kiese
322	Rottenbuch	Germany	48,175	11,642	750	8,4	1158	Temperate climate	Grassland	Ralf Kiese
323	Graswang	Germany	46,942	11,058	850	6,6	1359	Temperate climate	Grassland	Ralf Kiese
324,1	ES-SIC-BAR	Spain	40,780	-3,984	2170	9,0	599	Mediterranean climate	Alpine shrubland	Rosario G. Gavilán
324,2	ES-SIC-GUA	Spain	40,786	-3,978	2210	9,0	599	Mediterranean climate	Alpine grassland	Rosario G. Gavilán
324,3	ES-SIC-VAL	Spain	40,794	-3,961	2270	9,0	599	Mediterranean climate	Alpine grassland	Rosario G. Gavilán
324,4	ES-SIC-HEM	Spain	40,835	-3,966	2270	9,0	599	Mediterranean climate	Alpine grassland	Rosario G. Gavilán
325,01	IN-LAC, E-Ladakh/Changthang	India	33,008	78,416	5900	-7,8	250	Arid-temperate climate	Cold Himalyan Deserts, Subnival zone	Jiri Dolezal
325,02	IN-LAC, E-Ladakh/Changthang	India	32,980	78,363	5050	-3,5	150	Arid-temperate climate	Cold Himalyan Deserts, Alpine steppes	Jiri Dolezal
325,03	IN-LAC, E-Ladakh/Changthang	India	32,978	78,338	4720	-3	100	Arid-temperate climate	Cold Himalyan Deserts	Jiri Dolezal
329,01	IN-KJU-MGT	India	30,433	79,581	4254	3,1	1224	Subtropical arid climate	Alpine grassland	Sabyasachi Dasgupta
329,02	IN-KJU-GGT	India	30,456	79,580	3691	5,9	1472	Subtropical arid climate	Subalpin, Rhododendron scrub and grass land	Sabyasachi Dasgupta
330,00	US-PIO	USA	45,495	-112,483	2865	10	330	Temperate climate	Northern coniferous forest	Martha Apple
331,01	RO-CRO, SE Carpathians, Rodna Mts., Rebra Peak	Romania	47,585	24,635	2250	1,6	1255	Temperate climate	Alpine grassland	Mihai Pușcaș
331,02	RO-CRO, SE Carpathians, Rodna Mts., Buhăiescu Peak	Romania	47,582	24,632	2200	1,6	1255	Temperate climate	Alpine grassland	Mihai Pușcaș
331,03	RO-CRO, SE Carpathians, Rodna Mts., Gropile Peak	Romania	47,572	24,617	2050	1,6	1255	Temperate climate	Alpine grassland	Mihai Pușcaș
332,01	Vole (BLA_VOL)	Norway	61,896	9,143	1100	0,3	563	Boreal climate	Alpine Tundra (low alpine lichen heath)	Dirk Wundram
332,02	Derik (BLA_DER)	Norway	61,908	9,176	1221	-0,2	629	Boreal climate	Alpine Tundra (low alpine lichen heath)	Dirk Wundram
332,03	Skurvehøe (BLA_GRA)	Norway	61,895	9,221	1365	-0,5	713	Boreal climate	Alpine Tundra (low alpine lichen heath)	Dirk Wundram
332,04	Rundhøe (BLA_RUN)	Norway	61,905	9,246	1565	-1,1	804	Boreal climate	Alpine Tundra (mid alpine lichen heath)	Dirk Wundram
333	Patagonia	Argentina	-51,916	-70,407	165	6,4	202	Arid-temperate climate	Managed grassland	Pablo Peri
334,01	IT_ADO_GRM	Italy	46,331	11,563	2199	3,3	956	Temperate climate	Grassland	Brigitta Erschbamer
334,02	IT_ADO_PNL	Italy	46,383	11,593	2463	2	1118	Temperate climate	Grassland	Brigitta Erschbamer
334,03	IT_ADO_RNK	Italy	46,383	11,605	2757	0,8	1177	Temperate climate	Grassland & scree vegetation	Brigitta Erschbamer
334,04	IT_ADO_MTS	Italy	46,524	11,814	2893	-0,2	1121	Temperate climate	scree vegetation	Brigitta Erschbamer
335,01	IT_MAV_CCR	Italy	45,690	7,564	2340	3,8	1250	Temperate climate	Grassland with occasional larch	Umberto Morra di Cella
335,02	IT_MAV_LBA	Italy	45,640	7,550	2584	1,5	1250	Temperate climate	Alpine grassland	Umberto Morra di Cella
335,03	IT_MAV_PPE	Italy	45,650	7,540	2790	1,7	1250	Temperate climate	Scree vegetation	Umberto Morra di Cella
335,04	IT_MAV_CM	Italy	45,910	7,690	3014	-1,7	1200	Temperate climate	Scree vegetation	Umberto Morra di Cella
336	EC_ANT	Ecuador	-0,481	-78,141	5509	8	1336	Equatorial humid climate; Montane Grasslands and Shrublands	Grassland	Priscilla Muriel

(continued on next page)

Table 2s (continued)

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
337	EC_PIC	Ecuador	-0,177	-78,599	4676	10,6	1320	Equatorial humid climate; Montane Grasslands and Shrublands	Native grassland	Francisco Cuesta
338,01	IT_CAM_MAM	Italy	42,103	14,122	2722	2,9	898	Temperate climate	Alpine grassland	Angela Stanisci
338,02	IT_CAM_MAC	Italy	42,054	14,100	2625	2,9	898	Temperate climate	Alpine grassland	Angela Stanisci
338,03	IT_CAM_FEM	Italy	42,034	14,099	2411	2,9	898	Temperate climate	Alpine grassland	Angela Stanisci
339,01	EC_PNP1	Ecuador	-4,109	-79,162	3311	14,5	1163	Equatorial humid climate	Native shrubland	Marina Mazón
339,02	EC_PNP2	Ecuador	-4,106	-79,162	3352	14,5	1163	Equatorial humid climate	Native shrubland	Marina Mazón
339,03	EC_PNP3	Ecuador	-4,104	-79,161	3367	14,5	1163	Equatorial humid climate	Native shrubland	Marina Mazón
340,01	La Ly	Switzerland	46,031	7,249	2351	2,6	1544	Temperate climate	Dry subalpine-alpine grassland and heath, historical grazing but no more now	Jean-Paul Theurillat
340,02	Mt Brûlé	Switzerland	46,020	7,201	2547	2,6	1544	Temperate climate	Dry alpine grassland, no grazing	Jean-Paul Theurillat
341,01	IT -NAP-MOM	Italy	44,275	10,246	1842	5,7	1269	Temperate climate	Mosaic between primary subalpine shrublands and secondary grassland	Tomaselli Marcello
341,02	IT -NAP-CAS	Italy	44,331	10,207	1960	4,8	1055	Temperate climate	Subalpine secondary grassland	Tomaselli Marcello
341,03	IT -NAP-PCA	Italy	44,204	10,699	1803	5,1	992	Temperate climate	Subalpine secondary grassland	Tomaselli Marcello
341,04	IT -NAP-FOG	Italy	44,119	10,618	1696	5,1	1065	Temperate climate	Mosaic between primary subalpine shrublands and secondary grassland	Tomaselli Marcello
343,01	IN-KAS-GUL_1	India	34,015	74,206	3470	13,4	776	Temperate climate	Treeline of subalpine forest (dominated by <i>Betula utilis</i>)	Anzar A Khuroo
343,02	IN-KAS-GUL_2	India	34,015	74,205	3550	13,4	776	Temperate climate	Alpine scrub grassland (dominated by <i>Rhododendron-Juniperus</i>)	Anzar A Khuroo
343,03	IN-KAS-GUL_3	India	34,015	74,204	3640	13,4	776	Temperate climate	Alpine scrub grassland (dominated by <i>Rhododendron-Juniperus</i>)	Anzar A Khuroo
343,04	IN-KAS-GUL_4	India	34,015	74,204	3690	13,4	776	Temperate climate	Alpine scrub grassland (<i>Rhododendron-Juniperus</i> with Rock & Scree)	Anzar A Khuroo
344,1	ES-CPY-ACU	Spain	42,637	-0,062	2242	6,9	1383	Temperate climate	Subalpine environment	Juan J. Jiménez
344,2	ES-CPY-CUS	Spain	42,650	0,032	2519	4,9	1576	Temperate climate	Alpine (inferior)	Juan J. Jiménez
344,3	ES-CPY-TOB	Spain	42,656	-0,014	2779	4,9	1590	Temperate climate	Alpine	Juan J. Jiménez
344,4	ES-CPY-OLA	Spain	42,662	0,054	3022	3,4	1621	Temperate climate	Subnival rock	Juan J. Jiménez
346,01	BO-TUC (TucCop)	Bolivia	-16,224	-68,268	4862	4,0	785	Semi-arid tropical climate	Tropical dry alpine (Subnival), Grassland (Xerophytic Puna)	Rosa Isela Meneses
346,02	BO-TUC (TucPat)	Bolivia	-16,209	-68,270	5058	3,6	799	Semi-arid tropical climate	Tropical dry alpine (Nival), Grassland (Mesic Puna)	Rosa Isela Meneses
346,03	BO-TUC (TucWat)	Bolivia	-16,231	-68,258	4650	5,3	749	Semi-arid tropical climate	Tropical dry alpine (subnival), Grassland (Mesic Puna)	Rosa Isela Meneses
347,01	BO-SAJ (SajHui)	Bolivia	-18,118	-68,962	4567	2,5	382	Semi-arid tropical climate	Semi-arid tropical, climate (Subnival), Shrubland and grassland (Xerophytic Puna)	Rosa Isela Meneses
347,02	BO-SAJ (SarJas)	Bolivia	-18,155	-68,862	4931	4,3	373	Semi-arid tropical climate	Semi-arid tropical, climate (Nival), Shrubland and grassland (Xerophytic Puna)	Rosa Isela Meneses
347,03	BO-SAJ (SajPac)	Bolivia	-18,210	-68,968	4192	2,3	377	Semi-arid tropical climate	Semi-arid tropical climate (Alpin), Shrubland and grassland (Xerophytic Puna)	Rosa Isela Meneses
347,04	BO-SAJ	Bolivia	-18,127	-68,936	4759	5,6	344	Semi-arid tropical climate	Semi-arid tropical climate (Subnival), Shrubland and grassland (Xerophytic Puna)	Rosa Isela Meneses
348	Rostock-ECOLINK-Salix	Germany	54,061	12,085	13	8,5	590	Temperate climate	Willow short rotation coppice	Christel Baum
349	Kaltenborn (BIOTREE)	Germany	50,778	10,224	330	7,8	650	Temperate climate	Tree plantations	Michael Scherer-Lorenzen
351	Zedelgem (FORBIO)	Belgium	51,148	3,120	15	10,1	708	Temperate climate	Tree plantations	Kris Verheyen
352	Gedinne (FORBIO)	Belgium	49,986	4,981	397	10,4	670	Temperate climate	Tree plantations	Quentin Ponette
353	Hechtel-Eksel (FORBIO)	Belgium	51,165	5,313	56	8,6	1030	Temperate climate	Tree plantations	Bart Muys
354	Uppsala -ECOLINK-Salix	Sweden	60,439	18,080	22	5,6	470	Temperate climate	Arable Land	Martin Weih
355	Kreinitz	Germany	51,386	13,262	115	8,4	575	Temperate climate	Tree plantations	Anja Schmidt
356	IDENT-Macommer	Italy	13,817	8,700	640	13,8	866	Mediterranean climate	Abandoned fields in nursery	Simone Mereu
357	Bangor Diverse	UK	53,233	-4,133	10	9	1045	Temperate climate	NA	Andy Smith
358	MyDiv	Germany	51,392	11,886	115	8,8	507	Temperate climate	Agriculture	Olga Ferlian and Nico Eisenhauer
359	BEF-China Main Experiment: Site A	China	29,124	117,908	180	17,1	1777	Warm-temperate, humid climate	Subtropical broadleaf forest	Heike Feldhaar
360	Climate-match (Hucking, Kent, UK)	UK	53,397	-0,296	44	9,3	763	Temperate climate	Formerly Arable; Ungrazed pasture	Nadia Barsoum

361	ORPHEE	France	44,740	-0,797	60	12,75	876	Temperate climate	Pine plantation	Hervé Jactel
362	IDENT-Sault Ste. Marie	Canada	46,868	-84,571	210	-0,8	327	Temperate climate	Plantation	Bill Parker
363	IDENT-Montreal	Canada	45,858	-73,928	39	6,2	976	Temperate climate	High input agriculture	Alain Paquette
364	IDENT-Auclair	Canada	48,226	-69,100	333	2,3	1015	Temperate climate	Low input abandoned agriculture	Alain Paquette
365	IDENT-Cloquet	USA	46,679	-92,520	382	2,6	717	Temperate climate	Forest	Artur Stefanski
367	LTSERZAA_ORCHAMP_LORIAZ_1_LORI1370	France	46,031	6,925	1359	7,1	1207	Temperate climate	Mixed forest	Amélie Saillard
368	LTSERZAA_ORCHAMP_LORIAZ_2_LORI1620	France	46,033	6,921	1606	6	1170	Temperate climate	Coniferous forest	Amélie Saillard
369	LTSERZAA_ORCHAMP_LORIAZ_3_LORI1800	France	46,036	6,920	1785	6	1170	Temperate climate	Coniferous forest	Amélie Saillard
370	LTSERZAA_ORCHAMP_LORIAZ_4_LORI1930	France	46,036	6,915	1923	4,3	1104	Temperate climate	Forest-grassland ecotone	Amélie Saillard
371	LTSERZAA_ORCHAMP_LORIAZ_5_LORI2130	France	46,042	6,916	2125	2,7	975	Temperate climate	Subalpine grassland	Amélie Saillard
372	LTSERZAA_ORCHAMP_LORIAZ_6_LORI2330	France	46,045	6,913	2324	2,7	975	Temperate climate	Alpine meadow	Amélie Saillard
373	FR AME CBA - Cime des Barbarottes	France	44,303	6,937	2792	0,7	508	Temperate climate	Alpine scree	Philippe Choler
374	Hwange	Zimbabwe	-19,010	26,300	1010	21,6	524	Semi-arid tropical climate	Savannah	Hervé Fritz
393	ZAHG-2 Hwange National Park – Fixed vegetation plots	Zimbabwe	-19,010	26,500	1038	21,2	546	Semi-arid tropical climate	Savannah	Hervé Fritz
394	ZAHG-3 Hwange National Park – Sinamatella Mopane	Zimbabwe	-19,010	26,500	1038	21,2	546	Semi-arid tropical climate	Savannah	Hervé Fritz
395	ZAHG-4 Hwange National Park - Main Camp Waterhole transects	Zimbabwe	-19,010	26,500	1038	21,2	546	Semi-arid tropical climate	Savannah	Hervé Fritz
396	ZAHG-5 Magoli Village – Hwange District	Zimbabwe	-19,010	26,500	1038	21,2	546	Semi-arid tropical climate	Savannah	Hervé Fritz
397	Lamto	Ivory Coast	6,217	5,030	100	26,8	2146	Equatorial humid climate; tropical rain forest	Wet tropical savannah (transition tropical rain forest-Guinean savannah)	Jean-Christophe Lata
398	Comoé	Ivory Coast	9,115	-3,730	300	27,2	1096	Semi-arid tropical climate	West Sudanian savannah	Jean-Christophe Lata
399	Banco	Ivory Coast	5,394	-4,052	75	26,2	1738	Equatorial humid climate; tropical rain forest	Tropical rain forest	Jean-Christophe Lata
401	Kapiti	Kenya	-1,601	37,132	1646	17,8	1004	Semi-arid tropical climate	Rangeland	Victoria Carbonell
402	Nairobi	Kenya	-1,271	16,724	1857	18,9	592	Subtropical highland climate	Grassland	Victoria Carbonell
403	ZA Armorique-Pleine Fougères	France	48,488	-1,571	93	10,6	636	Temperate climate	Forest and wetland	Romain Georges
404	ZA Armorique - Sougeal	France	48,509	-1,512	70	10,6	636	Temperate climate	Wet grassland	Romain Georges
405	ZA Armorique - Rimou	France	48,398	-1,514	26	10,6	636	Temperate climate	Grassland	Romain Georges
406,01	SN1-MBU	Switzerland	46,641	10,238	2423	0,2	1143	Temperate climate	Grassland, rock and scree, no landuse	Sonja Wipf
406,02	SN1-MCH	Switzerland	46,644	10,234	2532	0,2	1143	Temperate climate	Grassland, rock and scree, no landuse	Sonja Wipf
406,03	SN1-CUO	Switzerland	46,716	10,171	2804	0,8	1146	Temperate climate	Nival rock and scree, no landuse	Sonja Wipf
407,01	SN2-MCS	Switzerland	46,736	10,428	2412	0,1	1179	Temperate climate	Grassland, rock and scree, low intensity cow grazing	Sonja Wipf
407,02	SN2-MIN	Switzerland	46,646	10,337	2507	0,4	1105	Temperate climate	Grassland, some low shrubs, some cow grazing	Sonja Wipf
407,03	SN2-MDG	Switzerland	46,694	10,331	2785	0,8	1146	Temperate climate	Grassland, rock and scree, low intensity cow grazing	Sonja Wipf
408	Lehavim LTER	Israel	31,359	34,847	460	18,7	318	Mediterranean climate	Rangeland	Marcelo Sternberg
409	Sde Boqer	Israel	30,868	34,772	475	18,8	90	Subtropical arid climate	Rangeland	Marcelo Sternberg
410	Kenting Karst Forest Dynamics Plot	Taiwan	21,966	120,816	260	24	2637	Equatorial humid climate	Natural tropical rain forest	Chiao-Ping Wang
411	Snowy Mountain_Mt Clarke	Australia	-36,420	148,280	2041	4,5	1979	Temperate climate	Alpine grassland	Ken Green
412	Garmisch	Germany	47,473	11,063	720	8	964	Temperate climate	Grassland	Ralf Kiese
413	Esterberg	Germany	47,517	11,158	1265	6,2	1043	Temperate climate	Grassland	Ralf Kiese
414	Dinderesso Forest	Burkina Faso	11,208	-4,403	397	27,1	1014	Semi-arid tropical climate	Savannah shrub	Jean-Christophe Lata
415	Chia-Yi Litchi Orchard	Taiwan	23,151	120,469	48	23,4	2338	Semi-arid tropical climate	Agriculture(Orchard)	Chi-Ling Chen
416	Gu-Keng Litchi Orchard	Taiwan	23,623	120,617	400	21,6	2637	Semi-arid tropical climate	Agriculture(Orchard)	Chi-Ling Chen
417	Min-Jian Tea Garden	Taiwan	23,817	120,651	413	22,6	2000	Semi-arid tropical climate	Agriculture(Tea Garden)	Chi-Ling Chen
418	Načetín	Czech Republic	50,590	13,254	775	5,4	789	Temperate climate	Spruce forest	Michal Ruzek
419	Načetín	Czech Republic	50,589	13,266	805	5,4	789	Temperate climate	Natural monocultural beech forest	Michal Ruzek
420,01	Toulouse-PYGAR-Auradé	France	43,559	1,064	157	12,4	730	Temperate climate	Agriculture (grass band along stream)	Jean-Luc Probst
420,02	Toulouse-PYGAR-Auradé	France	43,558	1,069	178	12,4	730	Temperate climate	Agriculture (fallow)	Jean-Luc Probst
420,03	Toulouse-PYGAR-Auradé	France	43,555	1,071	198	12,4	730	Temperate climate	Agriculture (grass fallow)	Jean-Luc Probst
421	Toulouse-PYGAR-Baget	France	42,955	1,031	522	9,3	964	Temperate climate	Grassland	Anne Probst
422,01	Toulouse-PYGAR-Bernadouze	France	42,803	1,424	1355	5,3	1191	Temperate climate	Peatland	Thierry Camboulive
422,02	Toulouse-PYGAR-Bernadouze	France	42,804	1,418	1433	7,2	952	Temperate climate	Forest	Thierry Camboulive
423	Toulouse-PYGAR-Météo	France	43,574	1,374	157	12,7	698	Temperate climate	Grassland	Christine Delire
424	Facundo	Argentina	-45,114	-69,987	460	9,3	162	Arid-temperate climate	Shrubland	Laura Yahdjian
425	Aldea beleiro	Argentina	-45,581	-71,392	640	5,9	497	Arid-temperate climate	Grasland	Laura Yahdjian

(continued on next page)

Table 2s (continued)

Site ID	Site	Country	Latitude	Longitude	Altitude (m asl)	MAT (°C)	MAP (mm)	Biome	Type of biotope	Contact
426	Rio Mayo	Argentina	-45,386	-70,253	460	9,2	192	Arid-temperate climate	Shrub-grass steppe	Laura Yahdjian
427	Las Chilcas	Argentina	-36,276	-58,266	12	15,1	930	Warm-temperate, humid climate	Grassland	Laura Yahdjian
428	Dinghushan	China	23,167	112,167	200–350	21,9	1773	Humid-arid tropical climate	NA	Jiangming Mo
429	Latnjajaure Climate change	Sweden	68,210	18,290	1000	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
430,01	Latnjajaure height transect 900–1400m	Sweden	68,210	18,290	900	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
430,02	Latnjajaure height transect 900–1400m	Sweden	68,210	18,290	1000	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
430,03	Latnjajaure height transect 900–1400m	Sweden	68,210	18,290	1100	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
430,04	Latnjajaure height transect 900–1400m	Sweden	68,210	18,290	1200	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
430,05	Latnjajaure height transect 900–1400m	Sweden	68,210	18,290	1300	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
430,06	Latnjajaure height transect 900–1400m	Sweden	68,210	18,290	1400	-2,7	659	Arctic climate	Alpine tundra	Juha Alatalo
431	Qatar 1 Acacia	Qatar	25,510	51,413	10	26,7	71	Subtropical arid climate	Acacia dryland	Juha Alatalo
432	Qatar 2 mangrove	Qatar	25,736	51,576	0	26,7	71	Subtropical arid climate	Acacia mangrove	Juha Alatalo
433	Qatar 3 Saltmarsh veg	Qatar	25,729	51,575	1	26,7	71	Subtropical arid climate	Arid saltmarsh with vegetation	Juha Alatalo
434	Qatar 4 mangrove planted	Qatar	25,661	51,548	0	26,7	71	Subtropical arid climate	Arid planted magrove	Juha Alatalo
435	Qatar 5 saltmarsh without veg	Qatar	25,657	51,544	1	26,7	71	Subtropical arid climate	Arid saltmarsh without vegetation	Juha Alatalo
436	Qatar 6 Grass	Qatar	25,221	51,294	10	26,7	71	Subtropical arid climate	Arid grassland	Juha Alatalo
437	Qatar 7 Zygophyllum	Qatar	25,233	51,294	10	26,7	71	Subtropical arid climate	Zygophyllum dryland	Juha Alatalo
438	Qatar 8 Acacia	Qatar	25,409	51,459	10	26,7	71	Subtropical arid climate	Acacia dryland	Juha Alatalo
439	Qatar 9 Mangrove	Qatar	25,697	51,550	0	26,7	71	Subtropical arid climate	Arid mangrove	Juha Alatalo
440	Qatar 10 saltmarsh veg	Qatar	25,698	51,552	1	26,7	71	Subtropical arid climate	Arid saltmarsh with vegetation	Juha Alatalo
440,01	E. Llebreta_ PN. Aiguestortes	Spain	42,921	1,480	1683	8,8	980	Temperate climate	Mountain grass	Esperança Gacia
440,02	Aiguadasi_ PN. Aiguestortes	Spain	42,954	1,552	1898	10,5	871	Temperate climate	Peatland forest	Esperança Gacia
440,03	Portarró_ PN. Aiguestortes	Spain	42,956	1,598	2046	10,5	871	Temperate climate	Mountain grass	Esperança Gacia
442	SERC GREW	USA	38,874	-76,547	1	12,9	1035	Temperate climate	Salt marsh	Thomas J. Mozdzer
443	PIE-LTER	USA	42,722	-70,847	2	9,5	1191	Temperate climate	Salt marsh	Thomas J. Mozdzer
444	Bylot Island	Canada	73,156	-79,972	20	-15,4	175	Arctic climate	Tundra	Vincent Maire
445	Umiujaq	Canada	56,552	-76,549	5	-5,4	525	Arctic climate	Tundra	Vincent Maire
446	Värriö	Finland	67,755	29,610	392	-1,3	537	Boreal climate	NA	Frank Berninger
447	Kahuzi-Biega	Democratic Republic Congo	-2,315	28,753	1900	14,9	1796	Equatorial humid climate; tropical rain forest	Natural forest (montane)	Marijn Bauters
448	Yoko	Democratic Republic Congo	0,295	25,302	400	24,9	1779	Equatorial humid climate; tropical rain forest	Natural forest (lowland)	Marijn Bauters
449	Yangambi Arboretum	Democratic Republic Congo	0,793	24,485	400	24,5	1770	Equatorial humid climate; tropical rain forest	Forest plantation	Marijn Bauters
451	Khibiny Station	Russia	67,637	33,725	320	-1,7	600	Boreal climate	Podsolich, peat	Yulia Zaika
452,01	Iskoras_Finnmark	Norway	69,417	25,612	350	-0,5	360	Boreal climate	tundra palsa mire (dry palsa w intact permafrost)	Casper T. Christiansen
452,02	Iskoras_Finnmark	Norway	69,417	25,612	350	-0,5	360	Boreal climate	tundra palsa mire (degrading palsa, degraded permafrost)	Casper T. Christiansen
452,03	Iskoras_Finnmark	Norway	69,417	25,612	350	-0,5	360	Boreal climate	tundra palsa mire (thaw pond, degraded permafrost)	Casper T. Christiansen
453,01	Galapagos WP169, Garrapatero - cinder cone	Ecuador	-0,698	-90,227	57	23,89	260	Subtropical arid climate	Semi-dry, deciduous vegetation	Heinke Jäger & Franz Zehetner
453,02	Galapagos WP171, Garrapatero - lava flow	Ecuador	-0,681	-90,225	47	23,91	276	Subtropical arid climate	Semi-dry, deciduous vegetation	Heinke Jäger & Franz Zehetner
453,03	Galapagos WP172, Garrapatero - cinder cone	Ecuador	-0,671	-90,248	210	22,99	302	Subtropical arid climate	Sub-tropical deciduous and evergreen shrubs and small trees	Heinke Jäger & Franz Zehetner
453,04	Galapagos WP180, Garrapatero - lava flow	Ecuador	-0,670	-90,254	231	22,62	315	Subtropical arid climate	Sub-tropical deciduous and evergreen shrubs and small trees	Heinke Jäger & Franz Zehetner
453,05	Galapagos WP174, Cerro Mesa - cinder cone	Ecuador	-0,642	-90,286	497	21,56	338	Subtropical arid climate	Mosaic of sub-tropical herb, evergreen shrub and tree vegetation, semi natural	Heinke Jäger & Franz Zehetner
453,06	Galapagos WP175, Cerro Mesa - lava flow	Ecuador	-0,642	-90,289	424	21,56	338	Subtropical arid climate	Mosaic of sub-tropical herb, evergreen shrub and tree vegetation, semi natural	Heinke Jäger & Franz Zehetner
453,07	Galapagos WP184, Cerro Crocker - cinder cone	Ecuador	-0,642	-90,326	866	19,87	398	Subtropical arid climate	Sub-tropical shrub and fern vegetation	Heinke Jäger & Franz Zehetner

453,08	Galapagos WP185, Cerro Crocker - lava flow	Ecuador	-0,645	-90,328	800	19,87	398	Subtropical arid climate	Sub-tropical shrub and fern vegetation	Heinke Jäger & Franz Zehetner
454	La Gamba	Costa Rica	8,700	-83,203	80	25,2	5748	Equatorial humid climate; tropical rain forest	Secondary forest	Florian Hofhansl
455	La Gamba	Costa Rica	8,705	-83,204	80	25,2	5748	Equatorial humid climate; tropical rain forest	Primary forest	Florian Hofhansl
456,01	Arctic station	Greenland	69,266	-53,464	89	-3	400	Arctic climate	Tundra	Regin Rønn
456,02	Arctic station	Greenland	69,269	-53,464	112	-3	400	Arctic climate	Tundra	Regin Rønn
457	FNQ Rainforest SuperSite, Daintree, Cape Tribulation (Rainforest)	Australia	-16,103	145,447	56	24,4	5143	Wet tropical rainforest	Natural rainforest	Michael Liddell
458	Tumbarumba Wet Eucalypt SuperSite	Australia	-35,657	148,152	1100	9,6	1274	Temperate climate	Managed wet eucalypt forest	Jacqui Stol
459	Warra Tall Eucalypt SuperSite	Australia	-43,099	146,684	100	10	1379	Temperate climate	Natural tall eucalypt forest	Timothy Wardlaw
460	Ayora	Spain	39,115	-0,950	1050	15,1	457	Mediterranean climate	Mediterranean mixed shrub	Alejandro Valdecantos
461	San Vicente Del Raspeig	Spain	38,384	-0,582	158	18	306	Mediterranean climate	Mediterranean mixed shrub	Alejandro Valdecantos
462	Albatera	Spain	38,230	-0,909	212	18,2	278	Mediterranean climate	Mediterranean mixed shrub	David Fuentes
463	Crevillente	Spain	38,244	-0,871	208	18,2	278	Mediterranean climate	Mediterranean mixed shrub	David Fuentes

References

- Adair, E.C., Parton, R.O.O.I.B.O.S.J., Del Grosso, S.J., Silver, W.L., Harmon, M.E., Hall, S.A., Hart, S.C., 2008. Simple three pool model accurately describes patterns of long-term litter decomposition in diverse climates. *Glob. Chang. Biol.* 14 (11), 2636–2660.
- Aerts, R., 1997. Climate, leaf litter chemistry and leaf litter decomposition in terrestrial ecosystems: a triangular relationship. *Oikos* 439–449.
- Aerts, R., 2006. The freezer defrosting: global warming and litter decomposition rates in cold biomes. *J. Ecol.* 94 (4), 713–724.
- Alsafran, M.H., Sarneel, J., Alatalo, J.M., 2017. Variation in Plant Litter Decomposition Rates Across Extreme Dry Environments in Qatar The Arab World Geographer. (In press).
- Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67 (1):1–48. <https://doi.org/10.18637/jss.v067.i01>.
- Berg, B., 2014. Decomposition patterns for foliar litter—a theory for influencing factors. *Soil Biol. Biochem.* 78, 222–232.
- Berg, B., McLaugherty, C., 2008. *Plant Litter: Decomposition, Humus Formation, Carbon Sequestration*. Springer Science & Business Media.
- Borer, E.T., Harpole, W.S., Adler, P.B., Lind, E.M., Orrock, J.L., Seabloom, E.W., Smith, M.D., 2014. Finding generality in ecology: a model for globally distributed experiments. *Methods Ecol. Evol.* 5 (1), 65–73.
- Bradford, M.A., Tordoff, G.M., Eggers, T., Jones, T.H., Newington, J.E., 2002. Microbiota, fauna, and mesh size interactions in litter decomposition. *Oikos* 99 (2), 317–323.
- Bradford, M.A., Li, R.J.W., Baldrian, P., Crowther, T.W., Maynard, D.S., Oldfield, E.E., King, J.R., 2014. Climate fails to predict wood decomposition at regional scales. *Nat. Clim. Chang.* 4 (7), 625.
- Bradford, M.A., Berg, B., Maynard, D.S., Wieder, W.R., Wood, S.A., 2016. Understanding the dominant controls on litter decomposition. *J. Ecol.* 104 (1), 229–238.
- Cornelissen, J.H., Van Bodegom, P.M., Aerts, R., Callaghan, T.V., Van Logtestijn, R.S., Alatalo, J., Hartley, A.E., 2007. Global negative vegetation feedback to climate warming responses of leaf litter decomposition rates in cold biomes. *Ecol. Lett.* 10 (7), 619–627.
- Cornwell, W.K., Cornelissen, J.H., Amatangelo, K., Dorrepaal, E., Eviner, V.T., Godoy, O., Qested, H.M., 2008. Plant species traits are the predominant control on litter decomposition rates within biomes worldwide. *Ecol. Lett.* 11 (10), 1065–1071.
- Cotrufo, M.E., Miller, M., Zeller, B., 2000. Litter decomposition. Carbon and Nitrogen Cycling in European Forest Ecosystems (pp. 276–296). Springer, Berlin, Heidelberg.
- Couteaux, M.M., Bottner, P., Berg, B., 1995. Litter decomposition, climate and litter quality. *Trends Ecol. Evol.* 10 (2), 63–66.
- Didion, M., Repo, A., Liski, J., Forsius, M., Bierbaumer, M., Djukic, I., 2016. Towards harmonizing leaves litter decomposition studies using standard tea bags—a field study and model application. *Forests* 7 (8), 167.
- Djukic, I., Zehetner, F., Watzinger, A., Horacek, M., Gerzabek, M.H., 2012. In situ carbon turnover dynamics and the role of soil microorganisms therein: a climate warming study in an Alpine ecosystem. *FEMS Microbiol. Ecol.* 83 (1), 112–124.
- Emmett, B.A., Beier, C., Estiarte, M., Tietema, A., Kristensen, H.L., Williams, D., ... Sowerby, A., 2004. The response of soil processes to climate change: results from manipulation studies of shrublands across an environmental gradient. *Ecosystems* 7 (6), 625–637.
- Fick, S.E., Hijmans, R.J., 2017. Worldclim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* 37 (12), 4302–4315.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Bala, G., 2006. Climate–carbon cycle feedback analysis: results from the C4MIP model intercomparison. *J. Clim.* 19 (14), 3337–3353.
- García Palacios, P., Maestre, F.T., Kattge, J., Wall, D.H., 2013. Climate and litter quality differentially modulate the effects of soil fauna on litter decomposition across biomes. *Ecol. Lett.* 16 (8), 1045–1053.
- García Palacios, P., Shaw, E.A., Wall, D.H., Hättenschwiler, S., 2016. Temporal dynamics of biotic and abiotic drivers of litter decomposition. *Ecol. Lett.* 19 (5), 554–563.
- Gavazov, K.S., 2010. Dynamics of alpine plant litter decomposition in a changing climate. *Plant Soil* 337 (1–2), 19–32.
- Gholz, H.L., Wedin, D.A., Smitherman, S.M., Harmon, M.E., Parton, W.J., 2000. Long-term dynamics of pine and hardwood litter in contrasting environments: toward a global model of decomposition. *Glob. Chang. Biol.* 6 (7), 751–765.
- Handa, I.T., Aerts, R., Berendse, F., Berg, M.P., Bröder, A., Butenschön, O., McKie, B.G., 2014. Consequences of biodiversity loss for litter decomposition across biomes. *Nature* 509 (7499), 218–221.
- Heim, A., Frey, B., 2004. Early stage litter decomposition rates for Swiss forests. *Biogeochemistry* 70, 299–313.
- Hoffman, G.E., Schadt, E.E., 2016. variancePartition: interpreting drivers of variation in complex gene expression studies. *BMC Bioinf.* 17.
- Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric models. *Biom. J.* 50 (3), 346–363.
- Houghton, R.A., 2007. Balancing the global carbon budget. *Annu. Rev. Earth Planet. Sci.* 35, 313–347.
- Ise, T., Moorcroft, P.R., 2006. The global-scale temperature and moisture dependencies of soil organic carbon decomposition: an analysis using a mechanistic decomposition model. *Biogeochemistry* 80 (3), 217–231.
- Johansson, M.B., Berg, B., Meentemeyer, V., 1995. Litter mass-loss rates in late stages of decomposition in a climatic transect of pine forests. Long-term decomposition in a Scots pine forest. *Can. J. Bot.* 73 (10), 1509–1521.
- Keuskamp, J.A., Dingemans, B.J., Lehtinen, T., Sarneel, J.M., Hefting, M.M., 2013. Tea bag index: a novel approach to collect uniform decomposition data across ecosystems. *Methods Ecol. Evol.* 4 (11), 1070–1075.
- Lal, R., 2004. Soil carbon sequestration impacts on global climate change and food security. *Science* 304 (5677), 1623–1627.
- Makkonen, M., Berg, M.P., Handa, I.T., Hättenschwiler, S., Ruijven, J., Bodegom, P.M., Aerts, R., 2012. Highly consistent effects of plant litter identity and functional traits on decomposition across a latitudinal gradient. *Ecol. Lett.* 15 (9), 1033–1041.
- McGuire, K.L., Treseder, K.K., 2010. Microbial communities and their relevance for ecosystem models: decomposition as a case study. *Soil Biol. Biochem.* 42 (4), 529–535.
- Mirtl, M., Borer, E., Burns, E., Djukic, I., Forsius, M., Haase, P., Haubold, H., Hugo, W., Jourdan, J., Lindemayer, D., McDowell, W.H., Muraoka, H., Orenstein, D., Peterseil, J., Shibata, H., Wohner, C., Yu, X., Pauw, J., 2018. Genesis, goals and achievements of long-term Ecological research at the global scale: a critical review of ILTER and future implications. *Sci. Total Environ.* (this issue).
- Moore, T.R., Trofymow, J.A., Prescott, C.E., Titus, B.D., CIDET Working Group, 2017. Can short-term litter-bag measurements predict long-term decomposition in northern forests? *Plant Soil* 1–8.
- Nadelhoffer, K.J., 2004. The DIRT experiment: litter and root influences on forest soil organic matter stocks and function. Chapter 15. In: Foster, D., Aber, J. (Eds.), *Synthesis Volume of the Harvard Forest LTER Program*. Oxford University Press, Oxford, UK.
- Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods Ecol. Evol.* 4, 133–142.
- Olson, J.S., 1963. Energy storage and the balance of producers and decomposers in ecological systems. *Ecology* 44 (2), 322–331.
- Parsons, S.A., Congdon, R.A., Lawler, I.R., 2014. Determinants of the pathways of litter chemical decomposition in a tropical region. *New Phytol.* 203 (3), 873–882.
- Parton, W., Silver, W.L., Burke, I.C., Grassens, L., Harmon, M.E., Currie, W.S., Fasth, B., 2007. Global-scale similarities in nitrogen release patterns during long-term decomposition. *Science* 315 (5810), 361–364.
- Pérez-Suárez, M., Arredondo-Moreno, J.T., Huber-Sannwald, E., 2012. Early stage of single and mixed leaf-litter decomposition in semiarid forest pine-oak: the role of rainfall and microsite. *Biogeochemistry* 108, 245–258.
- Persson, T., Van Oene, H., Harrison, A.F., Karlsson, P.S., Bauer, G.A., Cerny, J., ... Matteucci, G., 2000. Experimental sites in the NIPHYC/CANIF project. Carbon and Nitrogen Cycling in European Forest Ecosystems. Springer, Berlin Heidelberg, pp. 14–46.
- Pouyat, R.V., Setälä, H., Szlavecz, K., Yesilonis, I.D., Cilliers, S., Hornung, E., ... Whitlow, T.H., 2017. Introducing GLUSEEN: a new open access and experimental network in urban soil ecology. *J. Urban Econ.* 3 (1).
- Prescott, C.E., 2010. Litter decomposition: what controls it and how can we alter it to sequester more carbon in forest soils? *Biogeochemistry* 101 (1–3), 133–149.
- Soil Survey Staff, 2004. Soil survey laboratory methods manual. Soil Survey Investigations Rep. 42. USDA – NRCS, Washington, DC, USA.
- Trofymow, J.A., CIDET Working Group, 1998. The Canadian Intersite Decomposition Experiment (CIDET): Project and Site Establishment Report. vol. 378.
- Tuomi, M., Thum, T., Järvinen, H., Fronzek, S., Berg, B., Harmon, M., Liski, J., 2009. Leaf litter decomposition—estimates of global variability based on Yasso07 model. *Ecol. Model.* 220 (23), 3362–3371.
- Verheyen, K., De Frenne, P., Baeten, L., Waller, D.M., Hédli, R., Perring, M.P., De Lombaerde, E., 2016. Combining biodiversity resurveys across regions to advance global change research. *Bioscience* 67 (1), 73–83.
- Wall, D.H., Bradford, M.A., St. John, M.G., Trofymow, J.A., Behan-Pelletier, V., Bignell, D.E., Dangerfield, J.M., Parton, W.J., Rusek, J., Voigt, W., Wolters, V., Gardel, H.Z., Ayuke, F.O., Bashford, R., Beljakova, O.I., Bohlen, P.J., Brauman, A., Flemming, S., Henschel, J.R., Johnson, D.L., Jones, T.H., Kovarova, M., Kranabetter, J.M., Kutny, L., Lin, K.-C., Maryati, M., Masse, D., Pokarzhevskii, A., Rahman, H., Sabará, M.G., Salamon, J.-A., Swift, M.J., Varela, A., Vasconcelos, H.L., White, D., Zou, X., 2008. Global decomposition experiment shows soil animal impacts on decomposition are climate-dependent. *Glob. Chang. Biol.* 14 (11), 2661–2677.
- Walter, H., Breckle, S.W., 1999. *Vegetation und Klimazonen*. 544. Ulmer, Stuttgart.
- Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H., Wall, D.H., 2004. Ecological linkages between aboveground and belowground biota. *Science* 304 (5677), 1629–1633.
- Wickings, K., Grandy, A.S., Reed, S.C., Cleveland, C.C., 2012. The origin of litter chemical complexity during decomposition. *Ecol. Lett.* 15 (10), 1180–1188.
- Zhang, D., Hui, D., Luo, Y., Zhou, G., 2008. Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors. *J. Plant Ecol.* 1 (2), 85–93.