

Stomatal and Leaf Morphology Response of European Beech (*Fagus sylvatica* L.) Provenances Transferred to Contrasting Climatic Conditions

Peter Petřík ^{1,*}, Anja Petek ^{1,†}, Alena Konôpková ¹, Michal Bosela ¹, Peter Fleischer ^{1,2}, Josef Frýdl ³ and Daniel Kurjak ¹

¹ Faculty of Forestry, Technical University in Zvolen, T. G. Masaryka 24, 960 53 Zvolen, Slovakia; anja.petek1@gmail.com (A.P.); alena.konopkova@tuzvo.sk (A.K.); ybosela@tuzvo.sk (M.B.); xfleischer@tuzvo.sk (P.F.); kurjak@tuzvo.sk (D.K.)

² Institute of Forest Ecology, Slovak Academy of Sciences, L. Štúra 2, 960 53 Zvolen, Slovakia

³ Forestry and Game Management Research Institute, Strnady 136, 252 02 Jíloviště, Czech Republic; frydl@vulhm.cz

* Correspondence: peterpetrik94@gmail.com; Tel.: +421-915-830-543

† Contributed equally to this work.

Received: 12 November 2020; Accepted: 16 December 2020; Published: 18 December 2020

Abstract: Climate change-induced elevated temperatures and drought are considered to be serious threats to forest ecosystems worldwide, negatively affecting tree growth and viability. We studied nine European beech (*Fagus sylvatica* L.) provenances located in two provenance trial plots with contrasting climates in Central Europe. Stomata play a vital role in the water balance of plants by regulating gaseous exchanges between plants and the atmosphere. Therefore, to explain the possible adaptation and acclimation of provenances to climate conditions, stomatal (stomatal density, the length of guard cells, and the potential conductance index) and leaf morphological traits (leaf size, leaf dry weight and specific leaf area) were assessed. The phenotypic plasticity index was calculated from the variability of provenances' stomatal and leaf traits between the provenance plots. We assessed the impact of various climatic characteristics and derived indices (e.g., ecodistance) on intraspecific differences in stomatal and leaf traits. Provenances transferred to drier and warmer conditions acclimated through a decrease in stomatal density, the length of guard cells, potential conductance index, leaf size and leaf dry weight. The reduction in stomatal density and the potential conductance index was proportional to the degree of aridity difference between the climate of origin and conditions of the new site. Moreover, we found that the climate heterogeneity and latitude of the original provenance sites influence the phenotypic plasticity of provenances. Provenances from lower latitudes and less heterogeneous climates showed higher values of phenotypic plasticity. Furthermore, we observed a positive correlation between phenotypic plasticity and mortality in the arid plot but not in the more humid plot. Based on these impacts of the climate on stomatal and leaf traits of transferred provenances, we can improve the predictions of provenance reactions for future scenarios of global climate change.

Keywords: acclimation; adaptation; common garden; drought; ecodistance; mortality; phenotypic plasticity; stomatal frequency; stomatal size

1. Introduction

European beech forests may be seriously affected by climate change-induced drought due to their well-known vulnerability to water shortages [1,2]. Combinations of heat and drought stress may

cause a decrease in the vitality and competitive ability of beech populations [3–5]. There have been reports of beech populations facing strong selective pressures [6], which are foreseen to become more intense due to upcoming alterations in rainfall patterns and temperatures with ongoing climate change [7,8]. To mitigate these negative effects on the future performance of beech forests in afforestation programs in Europe, there has been increased interest in research on the intraspecific variation in beech responses to environmental changes [9–11].

Large intraspecific differences in morphological and physiological traits among the beech provenances of distinct origin reflect possible strategies which are expected to modify their response to drought. Beech populations show divergent water use strategies reflected in the differences of photosynthetic performance, water-use efficiency, leaf water potential, xylem embolism resistance and leaf morphology [10–13]. The intraspecific variation in tolerance of water deficit follows a pattern shaped by both regional and local scale effects. Beech populations originating from the sites with low precipitation [14–16], lower altitude [17] or marginal distribution range [9,18,19] show higher drought resistance in comparison with the populations from more humid environments. The observed functional variation between beech populations reaffirms the importance of local adaptation to water deficit in the context of climate change [13,20].

Common garden experiments allow us to assess the relative importance of adaptation to the site of origin and acclimation to the new environment in the expression of phenotypic traits, as all provenances are exposed to the same conditions in provenance trial plots [21]. As a result of adaptation to local original conditions, the performance and vitality of populations show a correlation with ecological characteristics at the site of origin, even after their transfer to a new environment. The effect of environmental change on a provenance planted at a given location can be expressed as the difference between the ecological characteristics of the trial plot and the site of provenance origin, called the ecodistance [22]. Moreover, if we study the performance of populations in different provenance trial plots and the differences between plots are greater than within, we expect that the differences between phenotypes are driven more by acclimation to current environmental conditions than by local adaptation [23,24]. Stomatal and leaf morphological traits such as stomatal density, potential conductance index and specific leaf area affect stomatal conductance and transpiration (functional traits) which in return influence performance, growth and survival [13,19]. Therefore, stomatal and leaf morphological traits represent a viable means to identify populations suitable for a specific environment.

The populations that possess stomatal and leaf morphological traits adapted to drought and heat stress will have an evolutionary advantage under future scenarios [3]. Hence, a plant strategy to cope with differences in water regimes involves altering stomatal density and stomatal size [25–27]. Some studies have shown that smaller stomata close more quickly than larger stomata do, thus indicating that this could enhance plant adaptation to drought [28]. However, there remains the debated issue of how stomatal density varies within a particular environment. It has been reported that drought resistant plants show higher stomatal density [27,29,30], but the results of more recent studies performed in controlled environments suggest that lower stomatal density improves drought tolerance [31,32]. Other functional traits frequently utilized in ecological studies and linked to drought tolerance are specific leaf area and leaf size. Several studies have revealed that changes in environmental factors such as light, temperature or nutrients strongly influence leaf traits [33,34]. Species with smaller, thicker leaves mainly occur in more stressful habitats and exhibit lower specific leaf areas. This trait is related to the species water use strategy [34], and it is highly plastic [35], although the precise physiological regulation mechanism of specific leaf area is still uncertain [33,36].

Furthermore, the phenotypic plasticity, defined as the capacity for a genotype to alter its morphology and/or physiology under altered environmental conditions [37], can play a major role in the survival and sustainability of forest populations subjected to global change [38–40]. This is generally seen as favourable under stress conditions because it enables plants to react to fluctuations in the environment [41,42]. However, several studies have reported a potential trade-off between phenotypic plasticity and individual fitness [43], suggesting reduced performance with increasing plasticity [38,44,45]. Therefore, it is crucial to assess the ability for stomatal and leaf morphological

traits to react plastically to their environment and to test the connection between their phenotypic plasticity and plant fitness and performance.

We investigated stomatal and leaf morphological traits, their phenotypic plasticity and a link to the climate of origin and current climate in nine European beech provenances located in two provenance trial plots with contrasting climates (warmer and drier/colder and more humid) in Central Europe. Based on the premise that environmental differences between provenance trial plots can alter the stomatal and leaf morphological traits of European beech, we hypothesized that (i) provenances that grow in drier and warmer provenance plots will exhibit lower values of measured traits than those growing in more humid and colder plots to increase their performance under suboptimal conditions. We further expected to find that (ii) the climate of the provenance's original site would affect the provenance's phenotype even 18 years after transfer to a different environment. In addition to the relationship between the climate of origin and phenotype itself, we also hypothesized that (iii) the provenance climate of origin should affect the phenotypic plasticity of provenances, where provenances from more heterogeneous environments show higher phenotypic plasticity. Finally, we hypothesized (iv) a negative relationship between phenotypic plasticity and tree mortality, and provenances with higher value of plasticity would acclimate better under different environments, thus mitigating the risk of mortality.

2. Materials and Methods

2.1. Locality Description and Plant Material

The material used for this experiment was collected from two European beech (*Fagus sylvatica* L.) provenance trial plots: Tále in the Slovak Republic (near Zvolen, 48°38'N, 19°02'E, 810 m a.s.l.) and Zbraslav in the Czech Republic (near Prague, 49°57'N, 14°22'E, 360 m a.s.l.). The Slovak provenance plot included loam soil with good nutrient availability and high water holding capacity, while the Czech provenance plot included sandy loam soil with poor nutrient availability and average water holding capacity [12]. Climate data for the original provenance sites were obtained from the WorldClim high-resolution climate database [46]. The climate characteristics of the Czech provenance plot were obtained from the Praha-Libuš meteorological station, and freely available data were provided by the Czech Hydrometeorological Institute (<http://portal.chmi.cz>). Climate data for the Slovak provenance plot were obtained from the nearby Kremnické Bane meteorological station monitored by the Slovak Hydrometeorological Institute. We calculated additional indices from the above climate data:

Ellenberg quotient (EQ) [47]

$$EQ = 1000 \times \frac{T_h}{Prec} \quad (1)$$

T_h —mean temperature of the hottest month (here July), $Prec$ —annual sum of precipitation.

Forest Aridity Index (FAI) [48]

$$FAI = 100 \times \frac{T_{7-8}}{Prec_{5-7} + Prec_{7-8}} \quad (2)$$

T_{7-8} —mean temperature of July and August, $Prec_{5-7}$ —precipitation sum for May to July, $Prec_{7-8}$ —precipitation sum for July to August.

Isothermality (ISO_T)

$$ISO_T = \frac{T \times (T_h - T_c)}{T_{MAX} - T_{MIN}} \times 100 \quad (3)$$

T —annual mean temperature, T_h —mean temperature of warmest month, T_c —mean temperature of coldest month, T_{MAX} —max temperature of warmest month, T_{MIN} —min temperature of coldest month.

Precipitation seasonality ($Seas_{Prec}$)

$$Seas_{prec} = \frac{\sigma_{Prec}}{Prec} \quad (4)$$

σ_{Prec} —annual standard deviation of precipitation, $Prec$ —annual sum of precipitation.

All geographical and climatic data are presented in Table 1. Optimal hydric conditions of European beech stands are represented by EQ values of below 20. European beech starts to lose its competitive performance in environments with EQ s of above 20 and is replaced by more xerotic tree species in places with EQ s of above 30 [49,50]. Moreover, locations with optimal rainfall patterns during the vegetation season for European beech are defined by an FAI of under 4.75 [48,51]. The EQ for the Czech provenance plot is 33.5, and the FAI is 5.1, which characterizes the plot as a location with marginal environmental conditions for European beech occurrence. The Slovak provenance plot with an EQ of 19.1 and FAI of 2.5 represents the optimal hydric environment according to the classifications mentioned above. Both provenance plots were established in 1998 as a part of the European provenance plot network, whereas provenances were planted as two-year-old seedlings [52]. Both plots were planted with 2×1 m spacing under a randomized block design with three blocks and fifty seedlings per block. Nine provenances were chosen for the analysis to capture the whole altitude range of the distribution of European beech (Figure 1). The distribution of provenance original sites regarding climate characteristics is visualized in Figure A1 of the Appendix. We were not able to sample some of the provenances in multiple blocks due to high spatial mortality. The third block of the Czech provenance plot completely died off, and some of the provenances remained only in one block. Similarly, the third block of the Slovak provenance plot suffered high mortality, and we avoided this to minimize sample heterogeneity due to unknown factors.

Table 1. Geographic and climate characteristics of provenance plots and their original locations.

| Provenance | Long | Lat | Alt | T | T ₅₉ | Prec | Prec ₅₉ | EQ | FAI | Isot | Seas _{prec} |
|-------------|-------|-------|------|------|-----------------|------|--------------------|------|-----|------|----------------------|
| FR04 | 2.58 | 44.15 | 850 | 10.8 | 16.8 | 804 | 344 | 23.8 | 5.4 | 3.8 | 14 |
| LUX12 | 6.2 | 49.67 | 400 | 8.6 | 14.9 | 866 | 365 | 19.7 | 4.8 | 3.1 | 11 |
| UK17 | −3.42 | 57.67 | 10 | 8.2 | 12.7 | 671 | 303 | 21.8 | 5.3 | 3.6 | 19 |
| SWE23 | 13.2 | 55.57 | 40 | 7.9 | 14.3 | 640 | 286 | 25.9 | 6.2 | 2.6 | 21 |
| GER26 | 10.67 | 53.65 | 55 | 8.3 | 15 | 678 | 319 | 25.5 | 5.1 | 3.0 | 17 |
| AU35 | 14.1 | 47.72 | 1250 | 2.4 | 9.2 | 1495 | 779 | 7.6 | 1.3 | 3.2 | 26 |
| AU36 | 14.85 | 47.53 | 1100 | 2.9 | 9.9 | 1168 | 648 | 10.4 | 1.6 | 3.1 | 32 |
| PL43 | 22.82 | 49.25 | 900 | 6.3 | 14.1 | 762 | 433 | 21.5 | 2.9 | 2.7 | 35 |
| PL67 | 18.17 | 54.33 | 250 | 5.8 | 13.2 | 633 | 336 | 24.6 | 4.3 | 2.4 | 30 |
| CZ Zbraslav | 14.37 | 49.95 | 360 | 8.25 | 15.6 | 532 | 330 | 33.5 | 5.1 | na | na |
| SK Tále | 19.03 | 48.63 | 850 | 6.58 | 14.1 | 842 | 441 | 19.1 | 2.5 | na | na |

Long—longitude, Lat—latitude, Alt—altitude, T—annual average temperature, T₅₉—average temperature during the vegetation season, Prec—sum of annual precipitation, Prec₅₉—sum of precipitation during the vegetation season, EQ—Ellenberg quotient, FAI—Forest Aridity Index, Isot—isothermality, Seas_{prec}—precipitation seasonality, na—not available.




Figure 1. Localities of the tested provenances (dots) and provenance plots (squares).

2.2. Stomatal and Leaf Morphological Traits

The samples were taken during June 2016 from full sun-exposed leaves located in the upper third of the crown to minimize irradiation and canopy position effects on stomatal and leaf morphology development [53,54]. We sampled six individuals per provenance per plot and made two imprints per individual. The imprints were made by the application of transparent nail polish to the abaxial side of the leaves [55]. The layer of polish was then transferred to a microscope slide with transparent tape. To avoid possible variations in stomatal distribution within the leaves, we took imprints between the second and third veins from the base of the leaves [17]. Six images were taken from each imprint using a Motic BA210 microscope with an integrated camera (Motic Electric, Linz, Austria). Three of these images were captured at 40×10 magnification, and three photos were captured at 10×10 magnification. The images at 10×10 magnification were used to assess stomatal density (SD). The number of stomata was calculated within a $750 \times 750 \mu\text{m}$ square per image with a random position using ImageJ 1.51k software (National Institute of Health, Bethesda, MD, USA). The assessed value of the number of stomata per square was converted to the number per square millimetre. The length of guard cells (L_A) was measured for ten stomata in a 40×10 magnification image using ImageJ software. The SD and L_A values were then averaged per individual for further analysis. Additional leaves from the same branch were scanned with a HP Scanjet G4010 scanner (Hewlett Packard, California, USA), and the leaf size (S_{leaf}) was subsequently measured by ImageJ software. The scanned leaves were stored in silica gel, after which the leaves were dried at 75°C to a constant weight (approximately 48 h). Afterwards, the dry weight (m_{leaf}) was assessed. From the measured parameters, we calculated the following traits:

The potential conductance index (PCI), an integrative variable of stomatal density and the length of guard cells, which can be used as a proxy for the theoretical maximal stomatal water vapour conductance [56,57]:

$$PCI = L_A^2 \times SD \times 10^{-4} \quad (5)$$

L_A —length of guard cells, SD —stomatal density.

Specific leaf area (SLA), a parameter that corresponds to the thickness and density of leaf lamina [58,59]:

$$SLA = \frac{S_{leaf}}{m_{leaf}} \quad (6)$$

S_{leaf} —leaf size and m_{leaf} —dry weight of leaves.

2.3. Quantification of Phenotypic Plasticity

The plasticity index based on maximum and minimum means (PI_v) was calculated for each trait and provenance, respectively [37].

$$PI_v = \frac{\bar{x}_{max} - \bar{x}_{min}}{\bar{x}_{max}} \quad (7)$$

\bar{x}_{max} —maximum mean (mean of the group showing the maximal value relative to that of other groups);
 \bar{x}_{min} —minimum mean (mean of the group showing the minimal value relative to that of other groups).
 In our case, the groups refer to identical provenances from two different provenance plots.

2.4. Statistical Analysis

Statistical analysis was performed using R statistical software (Version 4.0.3, R Core Team, Vienna, Austria). Original climate characteristics of provenances were analysed by principal component analysis to visualize likeness or disparity between the original sites of the provenances. The normal distribution of the obtained data was first tested by the Shapiro–Wilk test. A two-way analysis of variance was used where provenance and plots were set as factors with fixed effects. Moreover, differences between provenances were tested separately for each plot by Fisher’s LSD post hoc test. We used ecodistance to capture shifts between the climate of origin and the climate of the provenance plot. The ecodistance was defined as the difference between the investigated ecologically relevant variables at the test site and at the population origin [22,60]. Furthermore, mixed models in the R “nlme” package [61] were used to explore relationships between individual stomatal and leaf morphological traits (dependent variables) and various explanatory variables, such as *EQ*, *FAI*, latitude and longitude. To account for between-plot variability, we included plot as a random effect variable, and the between-plot variance in the intercept estimation was quantified. We were not able to include a block design in the mixed models, as there were missing data due to spatial mortality within the plots. We used maximum likelihood to estimate the parameters of the model. To estimate the importance of individual explanatory variables, we calculated the normalized model likelihoods (Akaike weights). First, we fitted all possible model variants, including the null model (including only the intercept) and full model (including all explanatory variables). Then, Akaike weights were calculated for each model based on the corrected Akaike information criterion using the “Weights” function in the “MuMIn” R package [62]. The models with the highest weights were further selected and interpreted. In addition to that, we calculated marginal R^2 (R^2_m) and conditional R^2 (R^2_c) for better comparison of fixed and random factors in model [63]. Statistical significance of differences between the plasticity of the traits was assessed by analysis of variance and Tukey’s post-hoc test. The relationships between original climate, phenotypic plasticity and tree mortality were tested by linear regression.

3. Results

3.1. Stomatal and Leaf Morphological Traits

The effects of the tested factors, provenance, plot and provenance-by-plot interactions were statistically significant for all traits except in the case of specific leaf area (*SLA*) (Table 2, Table A1 in Appendix A). Provenances growing in a warmer and drier site in the Czech Republic showed lower guard cell length (*L_A*, Figure 2A), stomatal density (*SD*, Figure 2B), potential conductance index (*PCI*, Figure 2C), leaf dry weight (*m_{leaf}*, Figure 2D) and leaf size (*S_{leaf}*, Figure 2E) values than the provenances in the colder and more humid Slovak plot. Provenances in the Czech provenance plot showed an average 57% reduction in the *PCI* for leaves that were 12% smaller than those in the Slovak provenance plot. Based on the average *m_{leaf}*, the provenances in the drier Czech plot accumulated 39% less biomass per leaf than the provenances in the more humid Slovak plot. Values of *SLA* were higher on average for the drier provenance plot, but changes were inconsistent and insignificant between provenances (Figure 2F).

Table 2. Results of two-way ANOVAs for each stomatal and leaf morphological trait.

| Factor | Df | Trait | L_A | SD | PCI | m_{leaf} | S_{leaf} | SLA |
|--------------------------|----|-------|--------|-------|---------|------------|------------|-------|
| Provenance | 1 | F | 57.95 | 17.65 | 11.59 | 3.79 | 10.83 | 0.69 |
| | | p | *** | *** | *** | ** | *** | 0.64 |
| Plot | 8 | F | 1771.5 | 457.4 | 1221.25 | 31.35 | 21.58 | 9.48 |
| | | p | *** | *** | *** | *** | *** | ** |
| Provenance \times Plot | 8 | F | 18.09 | 4.642 | 7.66 | 2.72 | 3.92 | 1.64 |
| | | p | *** | *** | *** | * | *** | 0.13 |

L_A —length of guard cells, SD —stomatal density, PCI —potential conductance index, S_{leaf} —leaf size, m_{leaf} —dry weight per leaf, SLA —specific leaf area, significance levels *** <0.001 ** <0.01 * <0.05.

3.2. Impact of Climate Ecodistance on Stomatal and Leaf Morphological Traits

The multifactorial approach based on mixed models showed that the models with singular explanatory variables performed better than those with multiple factors; thus, a further analysis employed simple linear regression models (Tables A2 and A3). The degree of aridity and temperature differences between provenances' origins and new plots (ecodistance) had a significant effect on provenances' stomatal development (Figure 3). The ecodistance of the Ellenberg quotient (EQ_{ED}) and forest aridity index (FAI_{ED}) had a significant negative influence on SD (Figure 3A,B) and the PCI (Figure 3C,D). Provenances transferred to a drier environment relative to their original site showed a proportionally lower density of stomata with lower potential conductance. The ecodistance of average temperature (T_{ED}) and average temperature during the vegetation season (T_{59ED}) also had a significant negative influence on SD , and provenances transferred to a climate warmer than that of their original site showed a proportional decrease in SD (Figure 3E,F). Other climate ecodistance indices showed no significant correlations with stomatal and leaf morphological traits.




Figure 2. Mean values and standard errors of stomatal and leaf morphological traits per provenance: length of guard cell (L_A , (A)), stomatal density (SD , (B)), potential conductance index (PCI , (C)), leaf size (S_{leaf} , (D)), leaf dry weight (m_{leaf} , (E)) and specific leaf area (SLA , (F)). The red colour represents provenances growing in the Czech provenance plot, and the blue colour represents provenances growing in the Slovak provenance plot. The dashed horizontal line represents the average per plot with the surrounding standard error interval band. Provenances are arranged based on the Ellenberg quotient of the original site (blue to red x axis band).




Figure 3. Linear regression results showing the relationship between ecodistance (EQ_{ED}, FAI_{ED}, T_{ED}, T_{59ED}) and stomatal morphological traits, stomatal density (SD, (A,B,E,F)) and the potential conductance index (PCI, (C,D)). Positive values on the horizontal axis represent transfer to a drier/hotter environment (red arrow), and negative values represent transfer to a more humid/colder environment (blue arrow).

3.3. Phenotypic Plasticity of Stomatal and Leaf Morphological Traits

To quantify the acclimation response, we calculated the phenotypic plasticity index for each trait and provenance. Plasticity indices of the potential conductance index (PCI_{PI}), leaf dry weight ($m_{leaf\ PI}$) and specific leaf area (SLA_{PI}) showed the highest values of plasticity among the provenances. In contrast, the plasticity index of the length of guard cells ($L_A\ PI$) was the lowest among the provenances (Figure 4). The climate of the original provenance locations affected the plasticity response of acclimation after transfer to a new environment. Provenances from locations with more heterogeneous environmental temperatures showed lower PCI_{PI} and $L_A\ PI$ plasticity. We found a positive relationship between the isothermality ($Isot$) of the original location and provenances PCI_{PI} and $L_A\ PI$ (Figure 5A,C). Moreover, provenances from environments with more heterogeneous precipitation distributions showed lower plasticity of L_A . The seasonality of precipitation ($Seas_{Prec}$) at the original locations of the provenances negatively influenced the $L_A\ PI$ of the provenances (Figure 5D). Furthermore, the original latitude (Lat) affected the PCI_{PI} , as provenances from lower latitudes showed a higher PCI_{PI} (Figure 5B). We found a significant negative relationship between provenance mortality and the plasticity of stomatal and leaf morphological traits (PCI_{PI} and SLA_{PI}) in the drier and hotter Czech provenance plot (Figure 6) but no significant relationship between mortality and plasticity in the more humid Slovak provenance plot.




Figure 4. Phenotypic plasticity indices of stomatal density (SD_{PI}), guard cell length ($L_A\ PI$), the potential conductance index (PCI_{PI}), leaf size ($S_{leaf\ PI}$), leaf dry weight ($m_{leaf\ PI}$) and specific leaf area (SLA_{PI}) for each provenance. Plasticity indices obtain values of 0 to 1, where 0 denotes no plasticity and 1 denotes theoretical maximal plasticity. Confidence intervals represent standard error and the capital letters correspond to results of post-hoc analysis.




Figure 5. Linear regression results showing the relationship between isothermality (Iso_T , (A,C)), latitude (Lat , (B)) and precipitation seasonality ($Seas_{Prec}$, (D)) of the provenance's original site and plasticity indices ($L_{A PI}$, PCI_{PI}).




Figure 6. Visualization of the relationship between plasticity indices (PCI_{PI} , (A); SLA_{PI} , (B)) and the observed mortality of provenances in the Czech (red) and Slovak (blue) provenance plot. The linear regression was significant only for drier Czech provenance plot.

4. Discussion

4.1. Functional Aspects of the Adaptive Response

Our study demonstrates that European beech provenances adjusted their stomatal and leaf morphological traits in response to being transferred to a new environment. Provenances exhibited significantly lower SD , L_A , PCI , S_{leaf} and m_{leaf} levels in the drier Czech provenance plot than in the more humid Slovak provenance plot. There were no significant differences in SLA when we considered provenance–plot interactions. Lack of significant differences regarding SLA might be caused by low sample size and high variability of the trait. Nevertheless, we observed that provenances in more xeric plot showed higher overall SLA . This is in opposition to general consensus [13,36,64,65], but has already been observed in some studies [66,67]. The adjustment of stomatal and leaf development can be seen as an adaptive response to either suboptimal climatic conditions of the Czech plot or to favourable climatic conditions of the Slovak plot. It has been reported that plants might improve their drought tolerance and water use efficiency (WUE) by reducing SD [31,32,68] and L_A [28,69]. Both herbaceous plants and trees react to episodic drought and long-term xericity of the environment by decreasing SD [25,70] and developing smaller stomata with lower L_A [26,56,71–73]. Combined stomatal morphology (L_A) and the distribution of stomata on leaves (SD), represented as the PCI , might be seen as a proxy for structural constraints of maximal stomatal conductance. A reduction in the PCI under xeric conditions should then ultimately reduce stomatal conductance [74] and water loss, which can lead to improved WUE [73,75]. Acclimation through the development of smaller leaves (S_{leaf}) under xeric conditions leads to less water loss through transpiration [76] and higher WUE [77,78]. It should be mentioned that WUE is also influenced by photosynthetic capacity and not just stomata related traits [13]. Plants exposed to water deficit show a reduction in m_{leaf} [64,79]. The combination of lower S_{leaf} with m_{leaf} in drier site might be explained by trees' strategy to invest more in root biomass with the cost of lower leaf biomass and leaf size [80]. The above-mentioned traits, therefore, represent plants' adaptive mechanisms in mitigating drought stress [81]. As these stomatal and leaf morphological traits have a significant impact on plant performance under water stress, their adjustment is vital for plants to successfully acclimate under changing conditions due to either anthropogenic transport to new environments or accelerating global climate change.

On the other hand, an increase in SD and L_A can enhance photosynthetic capacity [82,83], which could enhance tree performance under optimal climatic conditions where the strongest selective pressure is competition [84]. Leaves with higher SD and larger stomata (L_A) show an increase in maximal stomatal conductance [85–87], which leads to higher biomass accumulation and growth [88]. A higher PCI increases the maximal limit of stomatal conductance, which might improve photosynthetic capacity [89]. Moreover, the PCI has also been found to be related to leaf hydraulic conductance [57], which has been correlated with photosynthesis rates across plant species [89,90]. Higher S_{leaf} values under favourable conditions lead to higher photosynthesis rates [91,92], which positively affect leaf biomass production with higher m_{leaf} values [93]. According to our results, provenances in the drier and hotter Czech plot might acclimatise to their environment by undertaking a water conservation strategy with decreasing SD , L_A , PCI , S_{leaf} and m_{leaf} values. Conversely, provenances growing on the more humid Slovak plot could utilize the development of larger leaves (S_{leaf} , m_{leaf}) with higher SD , L_A and PCI values to maximize photosynthetic activity and growth in a competitive environment. Despite the clear theoretical basis for why the provenances showed significantly different values of the tested traits, we did not find any significant relationship between the tested traits and mortality. To address our first hypothesis (i), we found the morphological response as expected, but there is no evidence that the alternation of stomatal and leaf morphological traits had a significant positive effect on provenance performance. To capture the drought resistance profile of provenances for practical application, we suggest analysing additional physiological and functional traits, such as WUE , cuticular conductance, xylem embolism resistance and the turgor loss point [10,13,94]. Stomatal and leaf morphological traits alone are not satisfactory to define which provenances would be favourable for hotter and more xeric conditions in the near future.

4.2. Climate Ecodistance as an Effective tool for Provenance Research

We found that the *SD* and *PCI* of provenances depend on the aridity ecodistance (EQ_{ED} and FAI_{ED}) and temperature ecodistance (T_{ED} and T_{59ED}). The ecodistance represents the climatic shift between the original provenance site and the provenance plot to which it is transferred. The connection between both the climate of origin and the current climate and stomatal phenotype might suggest strong coordination between genetic and environmental impacts on stomatal development. A previous provenance study showed strong significant relationships between temperature and ecodistance and the phenology, morphology and dendrometric traits of European beech provenances [22]. Aridity ecodistance (EQ) has been found to be a significant explanatory variable for the vitality [95] and growth [96] of European beech provenances. A study of four temperate tree species (*Fagus sylvatica*, *Picea abies* (L.) Karst., *Pinus sylvestris* L. and *Quercus petraea* (Matt.) Liebl.) has also shown a significant relationship between aridity ecodistance (the annual aridity index) and the growth of provenances after transfer to a new environment [97]. Our second hypothesis (ii) is confirmed by a significant relationship between *SD*, the *PCI* and ecodistance, which incorporates the climate of origin as well as the current climate of new plots. The ecodistance has not attracted much popularity since its first formulation [22], but it seems that this simple mathematical formulation of climate transfer could capture the physiological, morphological and growth reactions of tree provenances to their new environments. More studies have focused on additional tree species, and other aspects of tree phenotyping are needed to test whether ecodistance is robust enough to be useful and reliable for forestry applications. The ecodistance could then be used not only to explain the effect of climatic shifts caused by the spatial transfer of provenances but also to predict provenance reactions to temporal changes caused by accelerating anthropogenic global climate change.

4.3. Phenotypic Plasticity

Populations with higher phenotypic plasticity can adapt to higher environmental variability and thus can minimize the risk of mortality [65,98,99]. Our results suggest that the *PCI*, m_{leaf} and *SLA* are the most plastic, while *L_A* is the least plastic trait among provenances. The low plasticity of *L_A* might suggest higher genetic control of stomatal size relative to the other tested traits. Similar results of low plasticity for *L_A* and high plasticity for the *PCI* and leaf morphology were also observed in beech provenance studies [71,95]. Furthermore, we found a significant relationship between climate heterogeneity of the original site (ISO_T and $Seas_{Prec}$) and the plasticity of *L_A* and *PCI* ($L_{A\ PI}$ and PCI_{PI}). We expected populations that had evolved under more heterogeneous environments to favour higher phenotypic plasticity to quickly adjust their phenotype if needed [11,37,99,100]. In contrast to what we expected, the provenances from the most heterogeneous environment showed the lowest $L_{A\ PI}$ and PCI_{PI} values, so hypothesis (iii) cannot be confirmed. We also found a negative significant relationship between latitude and PCI_{PI} , which might be attributable to a reduction in genetic diversity from lower to higher latitudes after the recolonization of habitats after the last glacial maximum [101–103]. A harsher northern environment or higher competition at the distribution edge may create more selection pressure than elsewhere in the distribution range [44]. Therefore, populations that evolved under strong selection pressure might favour more efficient strong genetic control over high phenotypic plasticity within the population [104]. This could lead to trait canalization which might be translated to lower phenotypic plasticity [105–107].

We found a significant positive relationship between PCI_{PI} , SLA_{PI} and mortality values, but only in the drier and hotter Czech provenance plots. However, in this study with phenotyping data from only the remaining trees, we cannot conclude that the high plasticity of these two traits causes high provenance mortality as the plasticity was calculated from surviving individuals and thus might be biased. Higher mortality, which created more open canopies and less competition among the remaining trees, might have caused the higher values of phenotypic plasticity, as trees reacted to newly available canopy space. High phenotypic plasticity is generally seen as a favourable property of plants, trees or populations under global climate change [37,108,109]. We cannot confirm our hypothesis that (iv) there is a negative relationship between phenotypic plasticity and tree mortality. Despite this, we find it important to discuss phenotypic plasticity. Both recent and earlier studies

have shown that the high phenotypic plasticity of plants should not be universally seen as a positive attribute [43,45,110,111]. The plasticity cost might not be pronounced under normal conditions, but when plants are exposed to a highly stressful environment, the plasticity cost might outweigh the fitness gain [112,113]. The results of several studies suggest reduced performance with increasing phenotypic plasticity under stress [38,44]. Thus, phenotypic plasticity, measured by common metrics [37], should not be automatically interpreted as beneficial for plants under global climate change. The high phenotypic plasticity of populations exposed to severe environmental stress might be associated with increased mortality and reduced fitness.

5. Conclusions

European beech provenances have shown a high degree of both adaptation and acclimation after transfer to a new environment. The observed differences in stomatal morphological traits were linked to the long-term aridity and air temperature of both the original site and the current provenance plot. The heterogeneity of the original site's climate and latitude affected the phenotypic plasticity of stomatal traits. Higher phenotypic plasticity was associated with higher mortality under suboptimal conditions but not under favourable hydric conditions. Additional functional and physiological traits should be considered to evaluate the resistance or performance of European beech provenances, as we have not found any direct link between mortality and the tested stomatal and leaf morphological traits. Ecodistance can be considered as an easy to use and robust tool for analysing adaptive responses of tree provenances under global climate change. Studies of phenotypic plasticity should not interpret the positive effects of high plasticity without taking into consideration the performance, fitness or vitality of plants.

Author Contributions: D.K., J.F. and A.K. conceived the experimental design and coordinated the experiment; P.P., A.P., A.K., J.F. and D.K. sampled the plant material; A.P. and P.P. processed the samples and performed the measurements; M.B., P.P. and P.F. performed the statistical analyses; P.P. and A.P. wrote the first version of the manuscript, which was discussed and approved by all of the co-authors. All authors have read and agreed to the published version of the manuscript.

Funding: The study was supported by research grants from the Slovak Research and Development Agency APVV-18-0390, by the Slovak Grant Agency for Science no. VEGA 1/0535/20 and by the Ministry of Agriculture of the Czech Republic, institutional support MZE-RO0118. Climatic data provided by SHMÚ and ČHMÚ are greatly appreciated.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Results of Fisher LSD post-hoc test.

| SK | LA | SD | PCI | S _{leaf} | m _{leaf} | SLA |
|-------|----|------|-----|-------------------|-------------------|-----|
| FR04 | BC | AC | AB | B | A | AB |
| LUX12 | AB | B | A | AB | ABC | BC |
| UK17 | AB | AB | AB | AB | AB | ABC |
| SWE23 | AB | C | B | AB | ABCD | A |
| GER26 | A | ABC | AB | A | ABC | C |
| AU35 | A | ABC | AB | AB | BCD | A |
| AU36 | AB | ABC | AB | A | D | AB |
| PL43 | AB | AB | A | AB | A | BC |
| PL67 | C | AC | A | A | CD | AB |
| CZ | LA | SD | PCI | S _{leaf} | m _{leaf} | SLA |
| FR04 | C | ABCD | AB | B | D | A |
| LUX12 | C | AB | B | AB | ACD | A |
| UK17 | AB | ABCD | ACD | A | ABC | A |
| SWE23 | AB | D | CD | A | B | A |
| GER26 | B | CD | D | A | AB | A |
| AU35 | AB | ABC | AC | A | AB | A |
| AU36 | AB | A | AB | A | ABC | A |
| PL43 | AC | BCD | ACD | AB | ABCD | A |
| PL67 | AC | ABCD | AB | AB | CD | A |

SD—stomatal density, LA—length of guard cells, PCI—potential conductance index, SLA—specific leaf area, S_{leaf}—leaf size, EQ_{ED}—Ellenberg quotient ecodistance, FAI_{ED}—Forest Aridity Index ecodistance, Long_{ED}—longitude ecodistance, Lat_{ED}—latitude ecodistance.


Table A2. AICc based variable weights of mixed models. Green to red spectrum represents the explanatory power of individual parameters and their combinations. Full model consists of all three parameters: aridity (expressed as the EQ and FAI), longitude and latitude.

| Trait | EQ _{ED} | Long _{ED} | Lat _{ED} | EQ _{ED} +Long _{ED} | EQ _{ED} +Lat _{ED} | Long _{ED} +Lat _{ED} | Full Model |
|-------------------|-------------------|--------------------|-------------------|--|---------------------------------------|--|------------|
| SD | 0.704 | 0.014 | 0.007 | 0.125 | 0.131 | 0.002 | 0.016 |
| LA | 0.222 | 0.31 | 0.168 | 0.202 | 0.031 | 0.047 | 0.021 |
| PCI | 0.211 | 0.461 | 0.151 | 0.065 | 0.032 | 0.073 | 0.007 |
| SLA | 0.312 | 0.239 | 0.292 | 0.044 | 0.06 | 0.046 | 0.006 |
| S _{leaf} | 0.16 | 0.472 | 0.152 | 0.069 | 0.031 | 0.102 | 0.014 |
| | FAI _{ED} | Long _{ED} | Lat _{ED} | FAI _{ED} + Long _{ED} | FAI _{ED} + Lat _{ED} | Long _{ED} + Lat _{ED} | Full model |
| SD | 0.263 | 0.056 | 0.029 | 0.046 | 0.532 | 0.009 | 0.065 |
| LA | 0.286 | 0.284 | 0.154 | 0.163 | 0.049 | 0.043 | 0.021 |
| PCI | 0.177 | 0.482 | 0.158 | 0.068 | 0.031 | 0.076 | 0.008 |
| SLA | 0.316 | 0.241 | 0.296 | 0.047 | 0.048 | 0.047 | 0.005 |
| S _{leaf} | 0.149 | 0.345 | 0.111 | 0.273 | 0.021 | 0.075 | 0.027 |

SD—stomatal density, LA—length of guard cells, PCI—potential conductance index, SLA—specific leaf area, S_{leaf}—leaf size, EQ_{ED}—Ellenberg quotient ecodistance, FAI_{ED}—Forest Aridity Index ecodistance, Long_{ED}—longitude ecodistance, Lat_{ED}—latitude ecodistance.

Table A3. Values of the marginal and conditional R^2 of mixed models presented in Table A2.

| | <i>SD</i> | | <i>L_A</i> | | <i>PCI</i> | | <i>SLA</i> | | <i>S_{leaf}</i> | |
|---|-----------|---------|----------------------|---------|------------|---------|------------|---------|-------------------------|---------|
| | R^2_m | R^2_c | R^2_m | R^2_c | R^2_m | R^2_c | R^2_m | R^2_c | R^2_m | R^2_c |
| <i>EQ_{ED}</i> | 0.38 | 0.76 | 0.03 | 0.87 | 0.01 | 0.89 | 0.04 | 0.86 | 0.02 | 0.38 |
| <i>Long_{ED}</i> | 0.01 | 0.84 | 0.02 | 0.78 | 0.01 | 0.91 | 0.01 | 0.75 | 0.12 | 0.18 |
| <i>Lat_{ED}</i> | 0.00 | 0.84 | 0.00 | 0.78 | 0.00 | 0.91 | 0.01 | 0.77 | 0.03 | 0.22 |
| <i>EQ_{ED} + Long_{ED}</i> | 0.38 | 0.75 | 0.07 | 0.92 | 0.01 | 0.91 | 0.04 | 0.86 | 0.19 | 0.75 |
| <i>EQ_{ED} + Lat_{ED}</i> | 0.37 | 0.77 | 0.03 | 0.86 | 0.01 | 0.88 | 0.04 | 0.86 | 0.05 | 0.40 |
| <i>Long_{ED} + Lat_{ED}</i> | 0.02 | 0.84 | 0.02 | 0.78 | 0.01 | 0.91 | 0.02 | 0.77 | 0.15 | 0.29 |
| Full model | 0.37 | 0.76 | 0.07 | 0.91 | 0.01 | 0.91 | 0.04 | 0.86 | 0.21 | 0.77 |
| | <i>SD</i> | | <i>L_A</i> | | <i>PCI</i> | | <i>SLA</i> | | <i>S_{leaf}</i> | |
| | R^2_m | R^2_c | R^2_m | R^2_c | R^2_m | R^2_c | R^2_m | R^2_c | R^2_m | R^2_c |
| <i>FAI_{ED}</i> | 0.06 | 0.82 | 0.02 | 0.83 | 0.00 | 0.90 | 0.02 | 0.80 | 0.09 | 0.43 |
| <i>Long_{ED}</i> | 0.01 | 0.84 | 0.02 | 0.78 | 0.01 | 0.91 | 0.01 | 0.75 | 0.12 | 0.18 |
| <i>Lat_{ED}</i> | 0.00 | 0.84 | 0.00 | 0.78 | 0.00 | 0.91 | 0.01 | 0.77 | 0.03 | 0.22 |
| <i>FAI_{ED} + Long_{ED}</i> | 0.06 | 0.82 | 0.04 | 0.85 | 0.01 | 0.91 | 0.02 | 0.79 | 0.23 | 0.58 |
| <i>FAI_{ED} + Lat_{ED}</i> | 0.11 | 0.86 | 0.02 | 0.83 | 0.00 | 0.90 | 0.02 | 0.79 | 0.09 | 0.40 |
| <i>Long_{ED} + Lat_{ED}</i> | 0.02 | 0.84 | 0.02 | 0.78 | 0.01 | 0.91 | 0.02 | 0.77 | 0.15 | 0.29 |
| Full model | 0.11 | 0.86 | 0.05 | 0.85 | 0.01 | 0.90 | 0.02 | 0.78 | 0.23 | 0.56 |

**Figure A1.** The principal component analysis biplot of provenance's original climate.

References

1. Leuschner, C.; Backes, K.; Hertel, D.; Schipka, F.; Schmitt, U.; Terborg, O.; Runge, M. Drought responses at leaf, stem and fine root levels of competitive *Fagus sylvatica* L. and *Quercus petraea* (Matt.) Liebl. trees in dry and wet years. *For. Ecol. Manag.* **2001**, *149*, 33–46, doi:10.1016/s0378-1127(00)00543-0.
2. Bréda, N.; Huc, R.; Granier, A.; Dreyer, E. Temperate forest trees and stands under severe drought: A review of ecophysiological responses, adaptation processes and long-term consequences. *Ann. For. Sci.* **2006**, *63*, 625–644, doi:10.1051/forest:2006042.
3. Gessler, A.; Keitel, C.; Kreuzwieser, J.; Matyssek, R.; Seiler, W.; Rennenberg, H. Potential risks for European beech (*Fagus sylvatica* L.) in a changing climate. *Trees* **2006**, *21*, 1–11, doi:10.1007/s00468-006-0107-x.

4. Kramer, K.; Degen, B.; Buschbom, J.; Hickler, T.; Thuiller, W.; Sykes, M.T.; De Winter, W. Modelling exploration of the future of European beech (*Fagus sylvatica* L.) under climate change—Range, abundance, genetic diversity and adaptive response. *For. Ecol. Manag.* **2010**, *259*, 2213–2222, doi:10.1016/j.foreco.2009.12.023.
5. Hacket-Pain, A.J.; Cavin, L.; Friend, A.D.; Jump, A.S. Consistent limitation of growth by high temperature and low precipitation from range core to southern edge of European beech indicates widespread vulnerability to changing climate. *Eur. J. For. Res.* **2016**, *135*, 897–909, doi:10.1007/s10342-016-0982-7.
6. Piovesan, G.; Biondi, F.; Di Filippo, A.; Alessandrini, A.; Maugeri, M. Drought-driven growth reduction in old beech (*Fagus sylvatica* L.) forests of the central Apennines, Italy. *Glob. Chang. Biol.* **2008**, *14*, 1265–1281, doi:10.1111/j.1365-2486.2008.01570.x.
7. Charru, M.; Seynave, I.; Morneau, F.; Bontemps, J.-D. Recent changes in forest productivity: An analysis of national forest inventory data for common beech (*Fagus sylvatica* L.) in north-eastern France. *For. Ecol. Manag.* **2010**, *260*, 864–874, doi:10.1016/j.foreco.2010.06.005.
8. Rita, A.; Gentilesca, T.; Ripullone, F.; Todaro, L.; Borghetti, M. Differential climate–growth relationships in *Abies alba* Mill. and *Fagus sylvatica* L. in Mediterranean mountain forests. *Dendrochronologia* **2014**, *32*, 220–229, doi:10.1016/j.dendro.2014.04.001.
9. Rose, L.; Leuschner, C.; Köckemann, B.; Buschmann, H. Are marginal beech (*Fagus sylvatica* L.) provenances a source for drought tolerant ecotypes? *Eur. J. For. Res.* **2009**, *128*, 335–343, doi:10.1007/s10342-009-0268-4.
10. Schuldt, B.; Knutzen, F.; Delzon, S.; Jansen, S.; Müller-Haubold, H.; Burlett, R.; Clough, Y.; Leuschner, C. How adaptable is the hydraulic system of European beech in the face of climate change-related precipitation reduction? *New Phytol.* **2015**, *210*, 443–458, doi:10.1111/nph.13798.
11. Aranda, I.; Bahamonde, H.A.; Sánchez-Gómez, D. Intra-population variability in the drought response of a beech (*Fagus sylvatica* L.) population in the southwest of Europe. *Tree Physiol.* **2017**, *37*, 938–949, doi:10.1093/treephys/tpx058.
12. Robson, T.M.; BeechCOSTe52 Database Consortium; Garzón, M.B. Phenotypic trait variation measured on European genetic trials of *Fagus sylvatica* L. *Sci. Data* **2018**, *5*, 180149, doi:10.1038/sdata.2018.149.
13. Sánchez-Gómez, D.; Robson, T.M.; Gascó, A.; Gil-Pelegrín, E.; Aranda, I. Differences in the leaf functional traits of six beech (*Fagus sylvatica* L.) populations are reflected in their response to water limitation. *Environ. Exp. Bot.* **2013**, *87*, 110–119, doi:10.1016/j.envexpbot.2012.09.011.
14. García-Plazaola, J.I.; Becerril, J. Effects of drought on photoprotective mechanisms in European beech (*Fagus sylvatica* L.) seedlings from different provenances. *Trees* **2000**, *14*, 485–490, doi:10.1007/s004680000068.
15. Peuke, A.D.; Schraml, C.; Hartung, W.; Rennenberg, H. Identification of drought-sensitive beech ecotypes by physiological parameters. *New Phytol.* **2002**, *154*, 373–387, doi:10.1046/j.1469-8137.2002.00400.x.
16. Bolte, A.; Czajkowski, T.; Cocozza, C.; Tognetti, R.; De Miguel, M.; Pšidová, E.; Ditmarová, L.; Dinca, L.; Delzon, S.; Cochard, H.; et al. Desiccation and Mortality Dynamics in Seedlings of Different European Beech (*Fagus sylvatica* L.) Populations under Extreme Drought Conditions. *Front. Plant Sci.* **2016**, *7*, 751, doi:10.3389/fpls.2016.00751.
17. Bresson, C.C.; Vitasse, Y.; Kremer, A.; Delzon, S. To what extent is altitudinal variation of functional traits driven by genetic adaptation in European oak and beech? *Tree Physiol.* **2011**, *31*, 1164–1174, doi:10.1093/treephys/tpu084.
18. Dounavi, A.; Netzer, F.; Čelepirović, N.; Ivanković, M.; Burger, J.; Figueroa, A.; Schön, S.; Simon, J.; Cremer, E.; Fussi, B.; et al. Genetic and physiological differences of European beech provenances (*F. sylvatica* L.) exposed to drought stress. *For. Ecol. Manag.* **2016**, *361*, 226–236, doi:10.1016/j.foreco.2015.11.014.
19. Thiel, D.; Kreyling, J.; Backhaus, S.; Beierkuhnlein, C.; Buhk, C.; Egen, K.; Huber, G.; Konnert, M.; Nagy, L.; Jentsch, A. Different reactions of central and marginal provenances of *Fagus sylvatica* to experimental drought. *Eur. J. For. Res.* **2014**, *133*, 247–260, doi:10.1007/s10342-013-0750-x.
20. Aranda, I.; Cano, F.J.; Gascó, A.; Cochard, H.; Nardini, A.; Mancha, J.A.; López, R.; Sánchez-Gómez, D. Variation in photosynthetic performance and hydraulic architecture across European beech (*Fagus sylvatica* L.) populations supports the case for local adaptation to water stress. *Tree Physiol.* **2015**, *35*, 34–46, doi:10.1093/treephys/tpu101.
21. De Villemereuil, P.; Gaggiotti, O.E.; Mouterde, M.; Till-Bottraud, I. Common garden experiments in the genomic era: New perspectives and opportunities. *Heredity* **2016**, *116*, 249–254, doi:10.1038/hdy.2015.93.

22. Matyas, C. Climatic adaptation of trees: Rediscovering provenance tests. *Euphytica* **1996**, *92*, 45–54, doi:10.1007/bf00022827.
23. Konôpková, A.; Kurjak, D.; Kmeť, J.; Klumpp, R.; Longauer, R.; Ditmarová, L.; Gömöry, D. Differences in photochemistry and response to heat stress between silver fir (*Abies alba* Mill.) provenances. *Trees* **2017**, *32*, 73–86, doi:10.1007/s00468-017-1612-9.
24. Kurjak, D.; Konôpková, A.; Kmeť, J.; Macková, M.; Frýdl, J.; Zivcak, M.; Palmroth, S.; Ditmarová, L.; Gömöry, D. Variation in the performance and thermostability of photosystem II in European beech (*Fagus sylvatica* L.) provenances is influenced more by acclimation than by adaptation. *Eur. J. For. Res.* **2018**, *138*, 79–92, doi:10.1007/s10342-018-1155-7.
25. Lavilla, M.; Seral, A.; Murciano, A.; Molino, S.; De-La-Fuente-Redondo, P.; Galan, J.M.G.Y. Stomatal traits in Iberian populations of *Osmunda regalis* (Osmundaceae, Polypodiopsida) and its relationship with bioclimatic variables. *Acta Bot. Malacit.* **2018**, *42*, 5–13, doi:10.24310/actabotanicaabmabm.v42i2.3029.
26. Durand, M.; Brendel, O.; Buré, C.; Le Thiec, D. Changes in irradiance and vapour pressure deficit under drought induce distinct stomatal dynamics between glasshouse and field-grown poplars. *New Phytol.* **2020**, *227*, 392–406, doi:10.1111/nph.16525.
27. Gindel, I. Stomatal Number and Size as Related to Soil Moisture in Tree Xerophytes in Israel. *Ecology* **1969**, *50*, 263–267, doi:10.2307/1934854.
28. Drake, P.L.; Froend, R.H.; Franks, P.J. Smaller, faster stomata: Scaling of stomatal size, rate of response, and stomatal conductance. *J. Exp. Bot.* **2013**, *64*, 495–505, doi:10.1093/jxb/ers347.
29. McCree, K.J.; Davis, S.D. Effect of Water Stress and Temperature on Leaf Size and on Size and Number of Epidermal Cells in Grain Sorghum 1. *Crop. Sci.* **1974**, *14*, 751–755, doi:10.2135/cropsci1974.0011183x001400050041x.
30. Cutler, J.M.; Rains, D.W.; Loomis, R.S. The Importance of Cell Size in the Water Relations of Plants. *Physiol. Plant.* **1977**, *40*, 255–260, doi:10.1111/j.1399-3054.1977.tb04068.x.
31. Hepworth, C.; Doheny-Adams, T.; Hunt, L.; Cameron, D.D.; Gray, J.E. Manipulating stomatal density enhances drought tolerance without deleterious effect on nutrient uptake. *New Phytol.* **2015**, *208*, 336–341, doi:10.1111/nph.13598.
32. Caine, R.S.; Yin, X.; Sloan, J.; Harrison, E.L.; Mohammed, U.; Fulton, T.; Biswal, A.K.; Dionora, J.; Chater, C.C.; Coe, R.A.; et al. Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. *New Phytol.* **2018**, *221*, 371–384, doi:10.1111/nph.15344.
33. Poorter, H.; Niinemets, Ü.; Poorter, L.; Wright, I.J.; Villar, R. Causes and consequences of variation in leaf mass per area (LMA): A meta-analysis. *New Phytol.* **2009**, *182*, 565–588, doi:10.1111/j.1469-8137.2009.02830.x.
34. Wright, I.J.; Reich, P.B.; Westoby, M. Strategy shifts in leaf physiology, structure and nutrient content between species of high- and low-rainfall and high- and low-nutrient habitats. *Funct. Ecol.* **2001**, *15*, 423–434, doi:10.1046/j.0269-8463.2001.00542.x.
35. Wellstein, C.; Chelli, S.; Campetella, G.; Bartha, S.; Galiè, M.; Spada, F.; Canullo, R. Intraspecific phenotypic variability of plant functional traits in contrasting mountain grasslands habitats. *Biodivers. Conserv.* **2013**, *22*, 2353–2374, doi:10.1007/s10531-013-0484-6.
36. Wellstein, C.; Poschlod, P.; Gohlke, A.; Chelli, S.; Campetella, G.; Rosbakh, S.; Canullo, R.; Kreyling, J.; Jentsch, A.; Beierkuhnlein, C. Effects of extreme drought on specific leaf area of grassland species: A meta-analysis of experimental studies in temperate and sub-Mediterranean systems. *Glob. Chang. Biol.* **2017**, *23*, 2473–2481, doi:10.1111/gcb.13662.
37. Valladares, F.; Sanchez-Gomez, D.; Zavala, M.A. Quantitative estimation of phenotypic plasticity: Bridging the gap between the evolutionary concept and its ecological applications. *J. Ecol.* **2006**, *94*, 1103–1116, doi:10.1111/j.1365-2745.2006.01176.x.
38. Merilä, J.; Hendry, A.P. Climate change, adaptation, and phenotypic plasticity: The problem and the evidence. *Evol. Appl.* **2014**, *7*, 1–14, doi:10.1111/eva.12137.
39. Stamp, M.A.; Hadfield, J.D. The relative importance of plasticity versus genetic differentiation in explaining between population differences; a meta-analysis. *Ecol. Lett.* **2020**, *23*, 1432–1441, doi:10.1111/ele.13565.
40. Palacio-López, K.; Beckage, B.; Scheiner, S.M.; Molofsky, J. The ubiquity of phenotypic plasticity in plants: A synthesis. *Ecol. Evol.* **2015**, *5*, 3389–3400, doi:10.1002/ece3.1603.
41. Blackman, C.J.; Aspinwall, M.J.; Tissue, D.T.; Rymer, P.D. Genetic adaptation and phenotypic plasticity contribute to greater leaf hydraulic tolerance in response to drought in warmer climates. *Tree Physiol.* **2017**, *37*, 583–592, doi:10.1093/treephys/tpx005.

42. Kramer, K.; Ducousso, A.; Gömöry, D.; Hansen, J.K.; Ionita, L.; Liesebach, M.; Lorent, A.; Schueler, S.; Sułkowska, M.; De Vries, S.; et al. Chilling and forcing requirements for foliage bud burst of European beech (*Fagus sylvatica* L.) differ between provenances and are phenotypically plastic. *Agric. For. Meteorol.* **2017**, *2017*, 172–181, doi:10.1016/j.agrformet.2016.12.002.
43. DeWitt, T.J.; Sih, A.; Wilson, D.S. Costs and limits of phenotypic plasticity. *Trends Ecol. Evol.* **1998**, *13*, 77–81, doi:10.1016/s0169-5347(97)01274-3.
44. Michalski, S.; Malyshev, A.V.; Kreyling, J. Trait variation in response to varying winter temperatures, diversity patterns and signatures of selection along the latitudinal distribution of the widespread grass *Lolium perenne* and plant *Arrhenatherum elatius*. *Ecol. Evol.* **2017**, *7*, 3268–3280, doi:10.1002/ece3.2936.
45. Kreyling, J.; Puechmaille, S.J.; Malyshev, A.V.; Valladares, F. Phenotypic plasticity closely linked to climate at origin and resulting in increased mortality under warming and frost stress in a common grass. *Ecol. Evol.* **2019**, *9*, 1344–1352, doi:10.1002/ece3.4848.
46. Fick, S.E.; Hijmans, R.J. WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* **2017**, *37*, 4302–4315, doi:10.1002/joc.5086.
47. Ellenberg, H. *Vegetation Ecology of Central Europe*; Cambridge University Press: Cambridge, UK; New York, NY, USA; New Rochelle, Australia; Melbourne, Australia; Sydney, Australia, 1988.
48. Führer, E.; Horváth, L.; Jagodics, A.; Machon, A.; Pödör, Z. Application of a new aridity index in Hungarian forestry practice. *Idojaras* **2011**, *3*, 205–216.
49. Fang, J.; Lechowicz, M.J. Climatic limits for the present distribution of beech (*Fagus* L.) species in the world. *J. Biogeogr.* **2006**, *33*, 1804–1819, doi:10.1111/j.1365-2699.2006.01533.x.
50. Stojanović, D.B.; Kržič, A.; Matović, B.; Orlović, S.; Duputić, A.; Djurdjević, V.; Galic, Z.; Stojnić, S. Prediction of the European beech (*Fagus sylvatica* L.) xeric limit using a regional climate model: An example from southeast Europe. *Agric. For. Meteorol.* **2013**, *176*, 94–103, doi:10.1016/j.agrformet.2013.03.009.
51. Gavrilov, M.B.; Lukić, T.; Janc, N.; Basarin, B.; Marković, S.B. Forestry Aridity Index in Vojvodina, North Serbia. *Open Geosci.* **2019**, *11*, 367–377, doi:10.1515/geo-2019-0029.
52. von Wühlisch, G.; Liesebach, M.; Hans, J. Muhs; Stephan, B.R. A network of international beech provenance trials. In *First EUFORGEN Meeting on Social Broadleaves*; Turok, J., Kremer, A., de Vries, S., Eds.; International Plant Genetic Resources Institute: Rome, Italy, 1998; pp. 164–172.
53. Vega, C.; González, G.; Bahamonde, H.A.; Valbuena-Carabaña, M.; Gil, L.; Fernández, V. Effect of irradiation and canopy position on anatomical and physiological features of *Fagus sylvatica* and *Quercus petraea* leaves. *Plant Physiol. Biochem.* **2020**, *152*, 232–242, doi:10.1016/j.plaphy.2020.05.007.
54. Aranda, I.; Pardo, F.; Gil, L.; Pardos, J. Anatomical basis of the change in leaf mass per area and nitrogen investment with relative irradiance within the canopy of eight temperate tree species. *Acta Oecologica* **2004**, *25*, 187–195, doi:10.1016/j.actao.2004.01.003.
55. Wolf, L. *Mikroskopická Tehnica*; National Health Publishing: Praha, Czech Republic, 1950.
56. Holland, N.; Richardson, A.D. Stomatal Length Correlates with Elevation of Growth in Four Temperate Species†. *J. Sustain. For.* **2009**, *28*, 63–73, doi:10.1080/10549810802626142.
57. Sack, L.; Cowan, P.D.; Jaikumar, N.; Holbrook, N.M. The “hydrology” of leaves: Co-ordination of structure and function in temperate woody species. *Plant, Cell Environ.* **2003**, *26*, 1343–1356, doi:10.1046/j.0016-8025.2003.01058.x.
58. Vile, D.; Garnier, É.; Shipley, B.; Laurent, G.; Navas, M.-L.; Roumet, C.; Lavorel, S.; Díaz, S.; Hodgson, J.G.; Lloret, F.; et al. Specific Leaf Area and Dry Matter Content Estimate Thickness in Laminar Leaves. *Ann. Bot.* **2005**, *96*, 1129–1136, doi:10.1093/aob/mci264.
59. Aranda, I.; Bergasa, L.F.; Gil, L. Effects of relative irradiance on the leaf structure of *Fagus sylvatica* L. seedlings planted in the understory of a *Pinus sylvestris* L. stand after thinning. *Ann. For. Sci.* **2001**, *58*, 673–680, doi:10.1051/forest:2001154.
60. Mátyás, C.; Fady, B.; Vendramin, G. Forests at the limit: Evolutionary—genetic consequences of environmental changes at the receding (xeric) edge of distribution. Report from a researcher workshop. *Acta Silv Lign Hung* **2009**, *66*, 201–204.
61. Pinheiro, J.; Bates, D.; DebRoy, S.; Sarkar, D.; EISPACK authors; Heisterkamp, S.; Van Willigen B.; Ranke J.; R Core Team. *nlme: Linear and Nonlinear Mixed Effects Models*; R package version 3.1-150; 2020. Available online: <https://cran.r-project.org> (accessed on 28 October 2020)
62. Bartoń, K. *MuMIn: Multi-Model Inference*; R package version 1.43.17; 2020. Available online: <https://cran.r-project.org> (accessed on 28 October 2020)

63. Nakagawa, S.; Schielzeth, H. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol. Evol.* **2012**, *4*, 133–142, doi:10.1111/j.2041-210x.2012.00261.x.
64. Marron, N.; Dreyer, E.; Boudouresque, E.; DeLay, D.; Petit, J.-M.; Delmotte, F.M.; Brignolas, F. Impact of successive drought and re-watering cycles on growth and specific leaf area of two *Populus x canadensis* (Moench) clones, “Dorskamp” and “Luisa_Avanzo”. *Tree Physiol.* **2003**, *23*, 1225–1235, doi:10.1093/treephys/23.18.1225.
65. Robson, T.M.; Sánchez-Gómez, D.; Cano, F.J.; Aranda, I. Variation in functional leaf traits among beech provenances during a Spanish summer reflects the differences in their origin. *Tree Genet. Genomes* **2012**, *8*, 1111–1121, doi:10.1007/s11295-012-0496-5.
66. Meier, I.C.; Leuschner, C. Genotypic variation and phenotypic plasticity in the drought response of fine roots of European beech. *Tree Physiol.* **2008**, *28*, 297–309, doi:10.1093/treephys/28.2.297.
67. Aspelmeier, S.; Leuschner, C. Genotypic variation in drought response of silver birch (*Betula pendula* Roth): Leaf and root morphology and carbon partitioning. *Trees* **2005**, *20*, 42–52, doi:10.1007/s00468-005-0011-9.
68. Dunn, J.; Hunt, L.; Afsharinafar, M.; Al Meselmani, M.; Mitchell, A.; Howells, R.; Wallington, E.; Fleming, A.J.; Gray, J.E. Reduced stomatal density in bread wheat leads to increased water-use efficiency. *J. Exp. Bot.* **2019**, *70*, 4737–4748, doi:10.1093/jxb/erz248.
69. Lawson, T.; Matthews, J. Guard Cell Metabolism and Stomatal Function. *Annu. Rev. Plant Biol.* **2020**, *71*, 273–302, doi:10.1146/annurev-arplant-050718-100251.
70. Bussotti, F.; Prancrazi, M.; Matteucci, G.; Gerosa, G. Leaf morphology and chemistry in *Fagus sylvatica* (beech) trees as affected by site factors and ozone: Results from CONECOFOR permanent monitoring plots in Italy. *Tree Physiol.* **2005**, *25*, 211–219, doi:10.1093/treephys/25.2.211.
71. Stojnić, S.; Orlovic, S.; Trudić, B.; Živković, U.; Von Wuehlisch, G.; Miljkovic, D. Phenotypic plasticity of European beech (*Fagus sylvatica* L.) stomatal features under water deficit assessed in provenance trial. *Dendrobiology* **2015**, *73*, 163–173, doi:10.12657/denbio.073.017.
72. Stojnić, S.; Orlović, S.; Miljković, D.; Galić, Z.; Kebert, M.; Von Wuehlisch, G. Provenance plasticity of European beech leaf traits under differing environmental conditions at two Serbian common garden sites. *Eur. J. For. Res.* **2015**, *134*, 1109–1125, doi:10.1007/s10342-015-0914-y.
73. Li, Y.; Li, H.; Li, Y.; Zhang, S. Improving water-use efficiency by decreasing stomatal conductance and transpiration rate to maintain higher ear photosynthetic rate in drought-resistant wheat. *Crop. J.* **2017**, *5*, 231–239, doi:10.1016/j.cj.2017.01.001.
74. Lawson, T.; Blatt, M.R.; Yu, L.; Shi, D.; Li, J.; Kong, Y.; Yu, Y.; Chai, G.; Hu, R.; Wang, J.; et al. Stomatal Size, Speed, and Responsiveness Impact on Photosynthesis and Water Use Efficiency. *Plant Physiol.* **2014**, *164*, 1556–1570, doi:10.1104/pp.114.237107.
75. Diazlopez, L.; Gimeno, V.; Simón, I.; Martínez, V.; Rodríguez-Ortega, W.; García-Sánchez, F. *Jatropha curcas* seedlings show a water conservation strategy under drought conditions based on decreasing leaf growth and stomatal conductance. *Agric. Water Manag.* **2012**, *105*, 48–56, doi:10.1016/j.agwat.2012.01.001.
76. George-Jaeggli, B.; Mortlock, M.Y.; Borrell, A.K. Bigger is not always better: Reducing leaf area helps stay-green sorghum use soil water more slowly. *Environ. Exp. Bot.* **2017**, *138*, 119–129, doi:10.1016/j.envexpbot.2017.03.002.
77. Anyia, A. Water-use efficiency, leaf area and leaf gas exchange of cowpeas under mid-season drought. *Eur. J. Agron.* **2004**, *20*, 327–339, doi:10.1016/s1161-0301(03)00038-8.
78. Songsri, P.; Jogloy, S.; Holbrook, C.; Kesmala, T.; Vorasoot, N.; Akkasaeng, C.; Patanothai, A. Association of root, specific leaf area and SPAD chlorophyll meter reading to water use efficiency of peanut under different available soil water. *Agric. Water Manag.* **2009**, *96*, 790–798, doi:10.1016/j.agwat.2008.10.009.
79. Poorter, L.; Markesteijn, L. Seedling Traits Determine Drought Tolerance of Tropical Tree Species. *Biotropica* **2008**, *40*, 321–331, doi:10.1111/j.1744-7429.2007.00380.x.
80. Slot, M.; Poorter, L. Diversity of Tropical Tree Seedling Responses to Drought. *Biotropica* **2007**, *39*, 683–690, doi:10.1111/j.1744-7429.2007.00328.x.
81. Basu, S.; Ramegowda, V.; Kumar, A.; Pereira, A. Plant adaptation to drought stress. *F1000Research* **2016**, *5*, 1554, doi:10.12688/f1000research.7678.1.
82. Tanaka, Y.; Sugano, S.S.; Shimada, T.; Hara-Nishimura, I. Enhancement of leaf photosynthetic capacity through increased stomatal density in *Arabidopsis*. *New Phytol.* **2013**, *198*, 757–764, doi:10.1111/nph.12186.

83. Harrison, E.L.; Cubas, L.A.; Gray, J.E.; Hepworth, C. The influence of stomatal morphology and distribution on photosynthetic gas exchange. *Plant J.* **2020**, *101*, 768–779, doi:10.1111/tpj.14560.
84. Davis, M.A.; Wrage, K.J.; Reich, P.B.; Tjoelker, M.G.; Schaeffer, T.; Muermann, C. Survival, growth, and photosynthesis of tree seedlings competing with herbaceous vegetation along a water-light-nitrogen gradient. *Plant Ecol.* **1999**, *145*, 341–350, doi:10.1023/a:1009802211896.
85. Franks, P.J.; Beerling, D.J. Maximum leaf conductance driven by CO₂ effects on stomatal size and density over geologic time. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 10343–10347, doi:10.1073/pnas.0904209106.
86. Fanourakis, D.; Briese, C.; Max, J.; Kleinen, S.; Putz, A.; Fiorani, F.; Ulbrich, A.; Schurr, U. Rapid determination of leaf area and plant height by using light curtain arrays in four species with contrasting shoot architecture. *Plant Methods* **2014**, *10*, 9, doi:10.1186/1746-4811-10-9.
87. Vráblová, M.; Vrábl, D.; Hronková, M.; Kubásek, J.; Šantrůček, J. Stomatal function, density and pattern, and CO₂ assimilation in *Arabidopsis thaliana* tmm1 and sdd1-1 mutants. *Plant Biol.* **2017**, *19*, 689–701, doi:10.1111/plb.12577.
88. Doheny-Adams, T.; Hunt, L.; Franks, P.J.; Beerling, D.J.; Gray, J.E. Genetic manipulation of stomatal density influences stomatal size, plant growth and tolerance to restricted water supply across a growth carbon dioxide gradient. *Philos. Trans. R. Soc. B Biol. Sci.* **2012**, *367*, 547–555, doi:10.1098/rstb.2011.0272.
89. Bucher, S.F.; Auerswald, K.; Tautenhahn, S.; Geiger, A.; Otto, J.; Müller, A.; Römermann, C. Inter- and intraspecific variation in stomatal pore area index along elevational gradients and its relation to leaf functional traits. *Plant Ecol.* **2016**, *217*, 229–240, doi:10.1007/s11258-016-0564-2.
90. Brodribb, T.J.; Holbrook, N.M.; Zwieniecki, M.A.; Palma, B. Leaf hydraulic capacity in ferns, conifers and angiosperms: Impacts on photosynthetic maxima. *New Phytol.* **2004**, *165*, 839–846, doi:10.1111/j.1469-8137.2004.01259.x.
91. Barigah, T.; Saugier, B.; Mousseau, M.; Guittet, J.; Ceulemans, R.J. Photosynthesis, leaf area and productivity of 5 poplar clones during their establishment year. *Ann. Sci. For.* **1994**, *51*, 613–625, doi:10.1051/forest:19940607.
92. Richards, R.A. Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp. Bot.* **2000**, *51*, 447–458, doi:10.1093/jexbot/51.suppl_1.447.
93. Gómez-del-Campo, M.; Ruiz, C.; Lissarrague, J.R. Effect of Water Stress on Leaf Area Development, Photosynthesis, and Productivity in Chardonnay and Airén Grapevines. *Am. J. Enol. Vitic.* **2002**, *53*, 138–143.
94. Wortemann, R.; Herbet, S.; Barigah, T.S.; Fumanal, B.; Alia, R.; Ducousso, A.; Gomory, D.; Roedel-Drevet, P.; Cochard, H. Genotypic variability and phenotypic plasticity of cavitation resistance in *Fagus sylvatica* L. across Europe. *Tree Physiol.* **2011**, *31*, 1175–1182, doi:10.1093/treephys/tptr101.
95. Čortan, D.; Nonić, M.; Šijačić-Nikolić, M. Phenotypic Plasticity of European Beech from International Provenance Trial in Serbia. In *Advances in Global Change Research*; Springer Science and Business Media LLC: Berlin/Heidelberg, Germany, 2019; pp. 333–351.
96. Petkova, K.; Molle, E.; Konner, M.; Knutzen, F. Comparing German and Bulgarian provenances of European beech (*Fagus sylvatica* L.) regarding survival, growth and ecodistance. *Silva Balc.* **2019**, *2*, 27–48.
97. Sáenz-Romero, C.; Kremer, A.; Nagy, L.; Újvári-Jármay, É.; Ducousso, A.; Kóczán-Horváth, A.; Hansen, J.K.; Mátyás, C. Common garden comparisons confirm inherited differences in sensitivity to climate change between forest tree species. *Peer J.* **2019**, *7*, e6213, doi:10.7717/peerj.6213.
98. Robson, T.M.; Rasztoivits, E.; Aphalo, P.J.; Alía, R.; Aranda, I. Flushing phenology and fitness of European beech (*Fagus sylvatica* L.) provenances from a trial in La Rioja, Spain, segregate according to their climate of origin. *Agric. For. Meteorol.* **2013**, *180*, 76–85, doi:10.1016/j.agrformet.2013.05.008.
99. Frank, A.; Pluess, A.R.; Howe, G.T.; Sperisen, C.; Heiri, C. Quantitative genetic differentiation and phenotypic plasticity of European beech in a heterogeneous landscape: Indications for past climate adaptation. *Perspect. Plant Ecol. Evol. Syst.* **2017**, *26*, 1–13, doi:10.1016/j.ppees.2017.02.001.
100. Gianoli, E.; Valladares, F. Studying phenotypic plasticity: The advantages of a broad approach. *Biol. J. Linn. Soc.* **2011**, *105*, 1–7, doi:10.1111/j.1095-8312.2011.01793.x.
101. Hewitt, G.M. Some genetic consequences of ice ages, and their role in divergence and speciation. *Biol. J. Linn. Soc.* **1996**, *58*, 247–276, doi:10.1006/bijl.1996.0035.
102. Hewitt, G.M. Post-glacial re-colonization of European biota. *Biol. J. Linn. Soc.* **1999**, *68*, 87–112, doi:10.1111/j.1095-8312.1999.tb01160.x.

103. Lander, T.A.; Klein, E.K.; Roig, A.; Oddou-Muratorio, S. Weak founder effects but significant spatial genetic imprint of recent contraction and expansion of European beech populations. *Heredity* **2020**, 1–14, doi:10.1038/s41437-020-00387-5.
104. Callahan, H.S.; Maughan, H.; Steiner, U.K. Phenotypic Plasticity, Costs of Phenotypes, and Costs of Plasticity. *Ann. N. Y. Acad. Sci.* **2008**, *1133*, 44–66, doi:10.1196/annals.1438.008.
105. McDonald, J.M.C.; Ghosh, S.M.; Gascoigne, S.J.L.; Shingleton, A.W. Plasticity Through Canalization: The Contrasting Effect of Temperature on Trait Size and Growth in *Drosophila*. *Front. Cell Dev. Biol.* **2018**, *6*, 156, doi:10.3389/fcell.2018.00156.
106. Debat, V.; David, P. Mapping phenotypes: Canalization, plasticity and developmental stability. *Trends Ecol. Evol.* **2001**, *16*, 555–561, doi:10.1016/s0169-5347(01)02266-2.
107. Klingenberg, C.P. Phenotypic Plasticity, Developmental Instability, and Robustness: The Concepts and How They Are Connected. *Front. Ecol. Evol.* **2019**, *7*, 7, doi:10.3389/fevo.2019.00056.
108. Yeh, P.J.; Price, T.D. Adaptive Phenotypic Plasticity and the Successful Colonization of a Novel Environment. *Am. Nat.* **2004**, *164*, 531–542, doi:10.1086/423825.
109. Nicotra, A.B.; Atkin, O.; Bonser, S.; Davidson, A.; Finnegan, E.; Mathesius, U.; Poot, P.; Purugganan, M.; Richards, C.; Valladares, F.; et al. Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.* **2010**, *15*, 684–692, doi:10.1016/j.tplants.2010.09.008.
110. Ghalambor, C.K.; McKay, J.K.; Carroll, S.P.; Reznick, D.N. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Funct. Ecol.* **2007**, *21*, 394–407, doi:10.1111/j.1365-2435.2007.01283.x.
111. Anderson, J.T.; Song, B. Plant adaptation to climate change—Where are we? *J. Syst. Evol.* **2020**, *58*, 533–545, doi:10.1111/jse.12649.
112. Van Buskirk, J.; Steiner, U.K. The fitness costs of developmental canalization and plasticity. *J. Evol. Biol.* **2009**, *22*, 852–860, doi:10.1111/j.1420-9101.2009.01685.x.
113. Scheiner, S.M. The genetics of phenotypic plasticity. XVI. Interactions among traits and the flow of information. *Evolution* **2018**, *72*, 2292–2307, doi:10.1111/evo.13601.

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).